This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

+ *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.

+ *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.

+ *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.

+ *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at `http://books.google.com/`

WILD BIRDS IN CITY PARKS

WALTER

Digitized by Google

HARVARD UNIVERSITY.

LIBRARY

OF THE

MUSEUM OF COMPARATIVE ZOÖLOGY

52,079

GIFT OF

George Howard Parker.

June 5, 1919.

With the Author's Compliments

JUN 5 1919

Digitized by Google

Digitized by Google

Wild Birds in City Parks

Being hints on identifying 200 birds, prepared primarily for the spring migration in Lincoln Park, Chicago, but adapted to other localities

BY

HERBERT EUGENE WALTER

AND

ALICE HALL WALTER

FOURTH ENLARGED REVISION
With Chart and Key

AUTHORS' EDITION
1910

Digitized by Google

Copyright 1901, 1902, 1904, 1910.
By Alice Hall Walter.

Digitized by Google

Contents.

	Page
Note to Fourth Revision	5
Preface	7
General Hints	9
Particular Hints	12
Water and Shore Birds	13
Grebes and Loons	13
Gulls and Terns	14
Ducks, Geese and Swans	15
Herons and Bitterns	19
Rails and Coots	20
Plover and Sandpipers	22
Land Birds	25
Bobwhite and Grouse	25
Doves	25
Birds of Prey	26
Cuckoos and Kingfishers	29
Woodpeckers	29
Swifts, Hummingbirds, etc	30
Perching Birds	31
Key to Land Birds	53
Table of Occurrence in Lincoln Park, Chicago	60
Table of Arrival in Lincoln Park, Chicago	64
Table of Distribution and Occurrence	71
Index to Common Names	89

Digitized by Google

"The first sparrow of spring! The year beginning with younger hope than ever! The faint silvery warblings heard over the partially bare and moist fields from the bluebird, the song sparrow and the red-wing, as if the last flakes of winter tinkled as they fell."

—*Thoreau's Walden.*

Digitized by Google

Note to the Fourth Edition.

A springtime has not yet passed without the arrival of the migrating birds. Upon this attractive subject an increasing number of new books appear every year, yet it seems probable from the way in which successive printings of this little book have been exhausted that there still exists a need for a simple, compact field manual at a minimum price.

The present edition is considerably enlarged by descriptions of over fifty additional birds, a revised key and a new table of comparative distribution. The entire subject matter also has undergone a thorough revision with the end in view of making the book useful in all localities in northeastern United States from the Mississippi to the Atlantic coast.

The authors wish to thank all those who have joined hands with them in the past and hope in the name of the birds that the circle may grow still larger in the days to come.

Digitized by Google

"All nature is so full that that district produces the greatest variety which is most examined."

—*Gilbert White*, 1768.

Digitized by Google

Preface.

The object of this little book is to furnish those who may be interested in making the acquaintance of wild birds with a simple letter of introduction to 200 birds, the majority of which are commonly seen during the spring migration.

Complete descriptions have been avoided, in the belief that the student should rely upon his own observation for the discovery of minor details. *The living bird* is the one important fact which will make the brief hints offered of value.

Anyone caring to make use of these hints may be assured that during the migrations of the birds city dwellers have one of the keenest delights of country life brought to their very doors, because many birds, migrating largely at night, are attracted by the lights of the city and stop off in their long journey to feed, so that a city park often contains a greater variety of feathered visitors than an equal area in the country.

We wish to remind those of our friends who have asked for pictures in a future edition that every springtime this book is copiously illustrated in our parks and around our homes by hundreds of living birds and that these illustrations are all life size, absolutely accurate in detail and colored true to nature.

Digitized by Google

"As for the birds * * * they add immeasurably to the wholesome beauty of life."

—*Theodore Roosevelt, in Bird Lore, Vol. II, p.* **98.**

Digitized by Google

General Hints.

"A good observer is quick to take a hint and follow it up."—*John Burroughs.*

The identification of birds depends quite as much upon accurate observation of their size, motions, flight, characteristic attitudes, manner of feeding, company, song, call-notes and haunts, as upon details of form and color. Especial care is necessary to insure correct estimates of size for the reason that living birds often appear smaller to the unpracticed eye than they actually are. The familiar *English Sparrow* is a convenient standard of size because it is usually at hand either in city or country for instant reference. Remembering that it is 6 *inches long* a practical though rough division of wild birds may be made as follows:—

1. Birds smaller than the *English Sparrow.*
2. Birds about the size of the *English Sparrow.*
3. Birds decidedly larger than the *English Sparrow.*

If a few general characteristics of the common bird-families be kept in mind, and these are quickly and almost unconsciously learned, the identity of a strange bird may usually be narrowed down to a few possibilities. For example:

Woodpeckers climb up and down the trunks of trees bracing with their tails and tapping the bark vigorously;

Nuthatches are smaller than woodpeckers and have much the same habit of climbing up and down tree-trunks but with a freer wig-wagging motion, often descending head downward;

Digitized by Google

Flycatchers sit erect with drooping tails, watching alertly for insect prey upon which they pounce in mid air, afterwards returning to their perch;

Swallows skim through the air in graceful and long sustained flights;

Sparrows have stout seed-cracking bills, feed upon the ground, seldom fly high or far at a time and are for the most part fine songsters;

Warblers are tiny, tireless, gaily-colored explorers of the twigs of trees and bushes;

Kinglets are smaller than warblers and quite as restless in their motions, but arrive earlier in the migration;

Wrens, with tails erect, slip mouse-like about brush heaps, crevices and bushes, though often perching in sight while singing;

Thrushes, who with the exception of the *Robin* and *Bluebird* are very plainly dressed, run about on the ground stopping suddenly in a listening attitude. When singing they fly up to some perch, although many of the unrivaled singers of this family are silent during their brief sojourn in city parks;

Vireos are most at home on the boughs of trees and sing freely as they glide in and out among the leaves to feed.

Female birds can usually be identified after the adult males have been seen, although the females and young of many species are obscurely marked or quite different from the adult males, a fact especially true of the warblers. Immature birds are not considered in the present scheme of

Digitized by Google

study since they are a source of confusion to the beginner and occur in any considerable number only during the fall migration.

The time of arrival, that is, when a bird may be expected during the migration, is a point worthy of particular attention. Many wild guesses may be avoided by simply noticing the dates when a bird has been known to arrive in any given locality from year to year.

By eliminating all birds whose arrival is decidedly earlier or later than that of the one in question, the matter is usually narrowed down to a few species whose identification is thus made much easier.

The safest way, however, is to make careful notes about a stranger on the spot. Memory is never more treacherous than in the case of the description of a doubtful bird when one appeals to some book of reference, a museum collection or the judgment of others. It is not advisable to attempt extensive "field-notes" at first but *it is essential* to keep a daily, dated list of all identified birds. A blank chart for this purpose is inserted in the back part of the book.

Never *chase* birds. Have patience, stand still a great deal and use your common sense all the time.

Do not make yourself believe that you see a certain bird because it has been reported by others. Do your own looking and listening and do it well.

If you cannot go birding alone take along as few friends as possible because birds are suspicious of human beings in flocks, especially when they move about much.

A pair of field or opera-glasses is a valuable aid, although practice is necessary in learning to adjust them rapidly and to fix them instantly upon the bird. Do not feel obliged to use the glasses when you are near enough to see well without them.

Keep the sun at your back, otherwise colors will deceive you.

Remember that birds do not stay "indoors" on account of rain, clouds or unfavorable weather. Warm spring rains literally fill the parks with feathered travelers who often tarry but a day.

Digitized by Google

Particular Hints.

For the convenience of those desiring access to more detailed sources of information the scientific name of each bird is placed in parenthesis below its common name, while prefixed to the common name is its number in the official list of the American Ornithologist's Union as revised to date.

Following the hints given about each bird are the names (in **black-faced type**) of any bird or birds for which it might easily be mistaken.

If information is desired as to the probable date of arrival of any bird mentioned, reference should be made to the table on page **65**.

The table beginning on page **74** indicates the birds which may be expected in given localities in northeastern United States.

Digitized by Google

Part 1.

WATER AND SHORE BIRDS.

3. HORNED GREBE. 14 in.

(Colymbus auritus.)

Sooty black, extending in narrow line up *back of neck*; rest of neck, sides and upper breast rich chestnut; glossy white below; head and chin black; buffy "horns" or crests, slanting backward from the eyes; slender black bill, tipped with yellow; eye, red; no tail. WINTER: black parts sooty; brown replaced by white; grayer below; crests and ruff less conspicuous. **Pied-billed Grebe.**

6. PIED-BILLED GREBE. 13½ in.

(Podilymbus podiceps.)

Brownish-black, showing gray on head and neck; belly, dusky white, otherwise brownish below; black throat-patch; a conspicuous black band across bill. WINTER: throat whitish; browner below; no band on bill. **Horned Grebe.**

7. LOON. 33 in.

(Gavia imber.)

Black, showing greenish on head and neck, spotted with square or oval patches of white on back and wings and striped with white in front of wings; white below; two conspicuous white-striped bars across the black neck giving the effect of a *broad black collar*; tail, very short; eye, red; large black bill, long and pointed; dives and sinks like the *Grebes*; note, an eerie, prolonged cry.

Digitized by Google

51. HERRING GULL. 24 in.

(Larus argentatus.)

White; wings and back, pearly blue-gray; bill mostly yellow; outer wing feathers partly black, marked and *tipped with white*; the closed wing shows black towards the end, marked with a regular line of white spots and tipped with white at the extreme point; looks large and heavy in flight. WINTER: Streaked on head and neck with gray. IMMATURE: Dark, much streaked with brownish; bill darker. **Ring-billed Gull. Kittiwake.**

54. RING-BILLED GULL. 19 in.

(Larus delawarensis.)

Similar to 51, but *smaller*; bill, greenish-yellow, bright at tip, *banded with black around the middle*; wing tipped at extreme point with black; feet greenish-yellow; eyelids ve-r milion-red. **Herring Gull.**

60. BONAPARTE GULL. 13 in.

(Larus philadelphia.)

Back and wings, light pearl-gray; head and throat, dark slate color; back of head, neck, underparts and square tail, white; wings tipped, and *bordered narrowly on the outside edge with black*; feet and legs, red; bill black. WINTER: hood, grayish white. **Common Tern.**

70. COMMON TERN. 15 in.

(Sterna hirundo.)

Pearly gray back and wings, whiter on rump; *top of head, shining black*; pure white on throat; dusky-white below; tail forked, *outer edge darkest*; bill long and red, blackening towards tip, held at right angles to the body in flight; feet reddish. WINTER: front of head and under-parts, white; bill nearly black. **Bonaparte Gull.**

Digitized by Google

129. AMERICAN MERGANSER. 25 in.

(Mergus americanus.)

Black; rump and tail, ashy; head, throat and upper part of neck, greenish; wing, *largely white*, edged with black and crossed by a black bar; white below, showing salmon tinge in flight; narrow white collar; long red bill tipped with black; eye, red. FEMALE: ashy-gray; *head and neck, brown;* salmon or brownish tinge across upper breast; *throat, white*; wings dark with a white patch; small crest on back of neck; feet orange; eye yellow. **American Golden-eye. Red-breasted Merganser.**

130. RED-BREASTED MERGANSER. 22 in.

(Mergus serrator.)

Long ragged crest; head and neck, black; broad white collar; broad cinnamon band streaked with black across breast, otherwise white below, showing salmon tinge in flight; wing largely white, edged and barred twice with black; eye and bill, red. FEMALE: back and wings dark gray turning to brown on head and neck; much paler on sides of neck and throat; gray across breast; crest small; white wing-patch. **American Merganser.**

131. HOODED MERGANSER. 18 in.

(Lophodytes cucullatus.)

Black, including throat and neck; *large circular crest, white bordered with black*; white below running up in front of the wings in two points; sides brownish, finely lined with black; white wing-patch, crossed by black bar; lengthwise white streaks on end of wings; *short* black bill; eye yellow. FEMALE: grayish-brown; throat white; crest small; sides unmarked. **Buffle-head.**

132. MALLARD. 23 in.

(Anas platyrhynchos.)

Gray-brown head, throat and neck glossy green to blue; *narrow* white collar; breast, rich brown; belly, gray lined with black; tail white, set off by black feathers curling up from either side of black rump; wing-patch purple, bordered on either side with a black and then a white bar. FEMALE: buffy-brown and black; much streaked and speckled; lighter on throat; shows wing-patch as in male.

Digitized by Google

133. BLACK DUCK. 22 in.

(Anas rubripes.)

Dark brown mottled with rusty, lighter below; wing-bar violet-purple edged with black; throat with few markings; legs brownish; bill greenish-black; *in flight white wing lining shows*; a suspicious pond and swamp duck often active at night; call, "quack". **Female Mallard.**

139. GREEN-WINGED TEAL. 14½ in.

(Nettion carolinensis.)

Gray, sides and shoulders finely lined with black; *white bar in front of wing*; head and throat brown with a broad green stripe from eye to back of head ending in a small tuft; wings, gray-brown with brilliant green and black wing-patch, buff-bordered; breast, pale chestnut speckled with black; buffy patches on sides of tail; white belly. FEMALE: mottled brown; top of head and back of neck dark brown; sides of head and neck buff, finely streaked; throat buff, unmarked; no green on head nor white bar in front of wings; wing-patch as in male but smaller. **Blue-winged Teal.**

140. BLUE-WINGED TEAL. 15 in.

(Querquedula discors.)

Back and underparts thickly mottled brownish and black, lightest below; head slaty with purplish gloss; *crescent-shaped white stripe in front of eye*; *shows blue* on bend of wing, followed by white bar and bright green wing-patch; white patch on sides of tail; bill black. FEMALE: dusky-brown; black on top of head; streaked and whitish on neck and sides of head; throat and about base of bill, *white*; back and underparts mottled and spotted; wing shows blue but no green; no white stripes on head. **Green-winged Teal.**

143. PINTAIL. 27 in.

(Dafila acuta.)

Head and throat brown, darkest on top; long, swan-like neck, black above, finely waved white and dusky on back and sides; *long middle tail-feathers*, black; white below with a conspicuous curved white stripe running up sides of neck to head and ending in a point; bronzy patch on wings. FEMALE: tail shorter but pointed; dusky, everywhere streaked; no white stripe on neck; whitish wing-bars; smaller than male. **Old-squaw.**

Digitized by Google

144. WOOD DUCK. 18 in.

(Aix sponsa.)

Highly variegated; long, glossy *crest* showing green and purple, marked by two narrow white parallel lines curving from bill and behind eye almost to end of crest; throat white extending irregularly in two stripes, one up behind eye, the other nearly around neck; breast glossy brown spotted with white and set off on either side by a conspicuous white stripe bordered with black; wings highly iridescent, marked by green patch bordered with white. FEMALE: duller, less conspicuously marked; head brownish-gray, slightly crested, greenish; throat and line extending from it around base of bill, with space about eye, white.

146. REDHEAD. 20 in.

(Marila americana.)

Puffy head and *upper* part of neck and throat, bright reddish-brown; breast and upper part of back with rump, *black*; belly white; middle back and *sides* evenly and finely waved black and white, *which shows gray at a distance*. FEMALE: grayish-brown, almost white on throat; head brown; lighter on neck. **Canvas-back.**

148. AMERICAN SCAUP DUCK. 18½ in.

(Marila marila.)

Similar to 149, but greenish on head; sides faintly barred; guttural call; less frequent on fresh water than 149. **Lesser Scaup.**

149. LESSER SCAUP DUCK. 16 in.

(Marila affinis.)

Head, neck and breast black, showing *purple* on head; back black and white, narrowly barred; upper part of wing dark, finely mottled with white; lower part of wing brownish-black with small white patch; lower breast and belly white, *sides barred;* bill, bluish tipped with black. FEMALE: black replaced by brown, lighter on head; white around bill. **Ring-necked Duck.** Female, easily confused with female **Lesser Scaup.**

Digitized by Google

151. AMERICAN GOLDEN-EYE. 19 in.

(Clangula clangula americana.)

Black, glossed with green; white below, extending around the neck in a collar; head bunchy, slightly crested and marked *below* and just back of eye by a nearly oval *white spot*; wings largely white; short bill. FEMALE: brown, snuff-colored on head; white collar and wing patch; band of bluish-gray across breast and down sides. **American Merganser.**

153. BUFFLE-HEAD. 15 in.

(Charitonetta albeola.)

Head, throat and upper neck, iridescent black; *black crest broadly banded with white over head* from eye to eye; back black, ashy on tail; wings black with large white patch; *collar and entire under parts, white.* FEMALE: smaller; brownish, lighter below; no crest; white patch on cheeks and small white wing-patch. **Hooded Merganser.**

154. OLD-SQUAW. 21 in.

(Harelda hyemalis)

WINTER PLUMAGE: Back, long tail-feathers and breast with *patch on sides of neck* black; throat, upper breast, top of head, neck and belly white; cheeks grayish-brown; wings black and gray; bill crossed by yellowish band. FEMALE: dark brown, lighter on head, grayish on tail; throat, breast and region about the eye gray-white; below white, brown on lower part of throat; no long tail-feathers; smaller than male. SUMMER PLUMAGE: head and neck, dusky black; grayish-white on sides of head; female shows more brown. **Pintail.**

165. WHITE-WINGED SCOTER. 19—20 in.

(Oidemia deglandi.)

Black, brownish below; *patch on wing* and *spot under eye, white* (nearly absent in winter); iris almost white; upper bill, red-orange on sides, *swollen on top*; fly high except in stormy weather. FEMALE: grayish-brown; whitish around upper bill and ears; white wing-patch; iris black. **American and Surf Scoters.**

Digitized by Google

167. RUDDY DUCK. 15—16 in.

(Erismatura jamaicensis.)

Upper half of head black, cheeks and chin white; elsewhere chestnut-red turning grayish; brown on rump and wings; belly white; tail blackish, *stiff* (often held erect by male while swimming); bill *broad*, short; feet large. WINTER PLUMAGE AND FEMALE: brown, somewhat mottled; *white stripe below eye;* swift, whirring flight; sink without diving or dive and swim below water like the grebes.

172. CANADA GOOSE. 40 in.

(Branta canadensis.)

Head and neck, *black with white throat-patch* extending up to and just behind eye; brownish-gray marked with lighter, darkest on back; rump and tail, black above and white below; fly after a leader in harrow-shaped ranks, often high up, crying "honk, honk, honk."

190. AMERICAN BITTERN. 30 in.

(Botaurus lentiginosus.)

Streaky, tawny and dark-brown, mixed with white and yellowish below; conspicuous black streak on sides of neck; short brown tail; *breast feathers loose and baggy*; yellow eye; call, loud resounding squawk, or booming notes which have given it the common name of "stake-driver."

191. LEAST BITTERN. 13 in.

(Ixobrychus exilis.)

Crown with back, tail and upper half of wings, shining greenish-black; rest of wing, buff and bright chestnut; throat and baggy breast, whitish shading into yellowish-brown on sides of neck and chestnut on back of neck; buffy-white below; tufts of dark feathers on either side of breast; yellow eye. **Green Heron.**

194. GREAT BLUE HERON. 45 in.

(Ardea herodias.)

Slaty-gray; top of head and *throat white*; *long* bill yellow beneath; legs *black*; *bend of wing reddish-brown;* blackish crest; feathers on breast long and streaky; stands on one leg; flies slowly with neck drawn in and legs extended behind; hoarse call. **Cranes.** (Cranes are rare; fly with necks extended; call, loud and resounding.)

Digitized by Google

201. GREEN HERON. 17 in.

(Butorides virescens.)

Wings and back dark green, latter tinged with blue-gray; crown greenish-black; neck and breast reddish-brown; throat and line down middle of neck to breast whitish, ending in light streaks on breast; the smallest heron; carries neck curved in flight; squawking note, "scow." **Night Heron.**

202. BLACK CROWNED NIGHT HERON. 24 in.

(Nycticorax nycticorax naevius.)

Green-black above; *sides of head*, throat and lower parts white; *ashy on lower half of wing* and tail; *neck short and thick*; pointed black bill; legs yellow; nests in flocks; feeds by night except in nesting season; call, "quawk." IMMATURE: brown, much streaked and mottled; appear slightly larger than adult. **Little Green Heron.**

212. VIRGINIA RAIL. 9½ in.

(Rallus virginianus.)

Streaked, blackish and olive-brown; back of neck and top of head plain brown; elsewhere *cinnamon-red* except *flanks* which are *heavily barred cross-wise*; long feet; short upturned tail; in fresh or salt marshes; grunting and guttural calls; one note of female much like sharp "click" of the Flicker. **King Rail. Clapper Rail.**

214. SORA. 8½ in.

(Porzana carolina.)

Olive-brown with lengthwise markings of black and some white; short tail turned up, showing buffy white underside; sides *barred* posteriorly with white; *front of head and throat black*; slaty line over eye; sides of neck and breast grayish-slate; bill *short and yellowish*; legs long; very stealthy; skulks and crouches in grass. YOUNG: no black; throat whitish, brown on breast. **Virginia Rail.**

Digitized by Google

219. FLORIDA GALLINULE. 13½ in.

(Gallinula galeata.)

Bluish-slate, washed with olive-brown on back, rump, wings and tail; grayish on belly with *white streaks along flanks*; under tail-coverts *white*; brilliant *red bill* and *plate* across forehead; legs green to red; call, "chuck" or clucking notes; walks and *swims*. **American Coot.**

221. AMERICAN COOT. 15 in.

(Fulica americana.)

Smooth blackish slate color, lighter below, black on head and neck; white bill shows *dark marks near the tip*; turned up tail; white underneath; wing edged with white; eye red; young show white below and no marks on bill; toes with scalloped edges. May be mistaken for a small duck. **Florida Gallinule.**

228. AMERICAN WOODCOCK. 10½ in.

(Philohela minor.)

Back, mixed black and rusty slate; below reddish-brown, no markings; *large* head with short neck and *very long straight bill*; eye set *high up and far back*; crown *barred crosswise* with black and rusty; short tail; burrows in soft mud for earthworms; nocturnal. **Wilson Snipe.**

230. WILSON SNIPE. 10½—11½ in.

(Gallinago delicata.)

Black and buff, much streaked; wings, dark brown marked with white; below white, streaked; sides and *lining of wings barred with black*; throat white; rusty tinge on breast and black-barred tail; *long, probing bill*; usually in fresh-water marshes or open swamps; *flight swift and crooked*; varied notes; call, harsh "scaipe"; makes a "bleating" sound (not vocal.) **Woodcock.**

Digitized by Google

239. PECTORAL SANDPIPER. 9 in.

(Pisobia maculata.)

Black mottled with rusty; white below, thickly streaked on breast; upper side of tail black, middle tail-feathers long and pointed; wet meadows; harsh creaky, whistling note; mating call musical, repeated eight times by male with throat greatly inflated. **Wilson Snipe. White-rumped Sandpiper.**

242. LEAST SANDPIPER. 5½—6 in.

(Pisobia minutilla.)

Blackish edged with reddish brown; light line over eye; shows whitish wing-bar; outer tail-feathers ash-gray; below white, finely streaked across breast; call sweet, "peep-peep"; in flocks along fresh and salt water; the *smallest* sandpiper. **Semipalmated Sandpiper.**

246. SEMIPALMATED SANDPIPER. 6 in.

(Ereunetes pusillus.)

Grayish-brown, marked with black and buffy; rump very dark; tail tapering; underparts *pure white*, slightly tinged and streaked across breast; white line over eye and dusky line beneath; note "peep-peep". **Other Sandpipers.**

248. SANDERLING. 8 in.

(Calidris leucophaea.)

SUMMER: mottled black, white and *rusty*; *white wing-bar*; below white, marked with black and tinged with *rusty* on throat and upper breast; *lacks the hind toe*; feet and bill black. WINTER: Grayish above, pure white below; call, a shrill whistle; mostly frequents beaches along the coast; runs rapidly, at times lifting its wings as if about to fly. **Spotted Sandpiper.**

254. GREATER YELLOW-LEGS. 15—16 in.

(Totanus melanoleucus.)

Black, much speckled with white; *head*, *neck* and *lower parts* white, streaked and marked with black except on middle of belly; bill *long*, *black*; *legs* and *feet yellow*; tail shows white barred with black; utters whistling call frequently; found near fresh and salt water. **Yellow-legs. Bartramian Sandpiper.**

Digitized by Google

255. YELLOW-LEGS. 11 in.

(Totanus flavipes.)

Similar in plumage and habits to 254, but *smaller* and more common; in flocks; most abundant in fall migration; call, three whistling notes resembling that of 254. **Greater Yellow-legs.**

256. SOLITARY SANDPIPER. 8½ in.

(Helodromas solitarius.)

Back, dusky olive-brown *finely spotted with white*, crown and back of neck showing dark streaks; white below, throat unmarked; sides of head and neck with breast slightly buffy and *distinctly streaked*; sides lightly barred; wings dark brown with one narrow white wing-bar; outer tail-feathers *white barred with black*; note, a soft whistle; frequents fresh water mostly. **Spotted Sandpiper.**

261. BARTRAMIAN SANDPIPER. 12 in.

(Bartramia longicauda.)

Black edged *regularly* with *buff*; neck *slender*, *buff*, *finely streaked*; tail *barred* with black; outer wing-feathers show *white* barring; below pale buff with arrow-marks of black; belly lighter with few markings; flute-like call or soft, mournful note; walks slowly, *nodding* head; flies high; when alighting *raises wings to full height before* folding them; grassy plains and pastures. **Golden Plover.**

263. SPOTTED SANDPIPER. 7½ in.

(Actitis macularia)

Greenish-ash marked lightly with black; *long white line over eye*; pure white below, *adults with dark round spots*; wings brownish, marked broadly with white bar; flies close to the water, wings full-spread, showing white bar; note, penetrating "pee-weet, weet"; walks with tilting motion. **Solitary Sandpiper.**

272. AMERICAN GOLDEN PLOVER. 10½ in.

(Charadrius dominicus.)

SUMMER: black, mottled with yellow; sides of breast white, otherwise black below; *forehead and line over eye*, *white*; below eye, *black*; lining of *wings ashy*. WINTER: whitish below. Plains and sandy hills or fields; runs swiftly; melodious whistle; in flocks. **Black-bellied Plover.**

Digitized by Google

273. KILLDEER. 10 in.

(Oxyechus vociferus.)

Gray-brown; rump rusty-brown; tail rather long; white below; white collar followed by a black collar; *black band across breast*; forehead, line over eye and wing-bar white; bill black; runs very swiftly; note, loud and persistent "killdee," often heard high overhead; found mostly inland.

274. SEMI-PALMATED PLOVER. 6¾ in.

(Aegialitis semipalmata.)

Forehead, throat and *collar around neck white, followed by a black collar* broadening across breast; below white; gray-brown above; in small scattering flocks along beaches or water-courses; call, very sweet, two or three notes, ascending; rather silent but not shy. **Piping Plover.**

Digitized by Google

Part 11.

LAND BIRDS.

289. BOB-WHITE. 10 in.

(Colinus virginianus.)

MALE: reddish-chestnut, much variegated with black and buff; tail ashy; *throat and line over eye white*; forehead, line below eye and band across breast, black; belly white, finely barred with wavy black lines; spring call of male, "bob-white"; varied notes; in flocks except during breeding season; *highly beneficial.* FEMALE: white markings buffy, little black on breast.

300. RUFFED GROUSE. 17 in.

(Bonasa umbellus.)

Black and reddish-buff, *much barred*; below barred with grayish-white; tail mottled showing broad black bar; *black feather-tufts* on side of neck (longest on male) which form a ruff; woodland and shrubby pastures; *call* of male, a remarkable whirring of the wings described as *"drumming;"* call of female "cluck" or a note of distress when young are disturbed. **Canada Grouse. Prairie Hen.**

316. MOURNING DOVE. 12 in.

(Zenaidura macroura carolinensis.)

Gray-blue, tinged with brownish; tail broadly tipped with white, *tapering to a narrow point*; sides of neck iridescent; marked with black on wings and sides of head; rapid whistling flight; note, "coo-oo." **Pigeon. Cuckoos.**

Digitized by Google

331. MARSH HAWK. 22 in.

(Circus hudsonius.)

Bluish-gray, showing light streaks; gray below, belly white marked with brown; tail barred; *white rump;* (female brown, barred and streaked more heavily); note shrill.

332. SHARP-SHINNED HAWK. 12 in.

(Accipiter velox.)

Slate-gray, crown darker; edge of wing barred with black; *tail long, square*, barred broadly, white-tipped; below white slightly tinged with brown, streaked or faintly barred; *usually silent* except at nesting period; soars in narrow circles; flies swiftly; attacks poultry. **Cooper Hawk.**

333. COOPER HAWK. 17 in.

(Accipiter cooperi.)

Resembles closely 332, but is *larger*, crown almost black; tail *rounded* and showing more white at tip; flies *swiftly*; attacks poultry; cackling note. **Sharp-shinned, Red-shouldered and Red-tailed Hawks.**

337. RED-TAILED HAWK. 19—22 in.

(Buteo borealis.)

Much like 339 above but without shoulder-patch; *tail rich, red-brown*, barred *once* with black (more or less completely barred in female and immature); below white; upper breast and throat streaked (*adults show red-brown tinge* on sides); call, a squealing whistle; woodland; beneficial. **Red-shouldered and Broad-winged Hawks.**

339. RED-SHOULDERED HAWK. 18—20 in.

(Buteo lineatus.)

Dark brown varied with reddish-brown, buffy or grayish; edge of wing barred black and white, showing *four notches* in *flight;* red-brown shoulder-patch; tail blackish, barred and tipped with white; below reddish-buff, *barred* with whitish; (immature *streaked* below or *heavily spotted*; shoulder-patch less showy); note, "kee-you"; beneficial, although commonly known as a hen-hawk. **Red-tailed and Broad-winged Hawks.**

Digitized by Google

343. BROADWINGED HAWK. 14—18 in.

(Buteo platypterus.)

Dark brown, slightly mottled with reddish-buff; tail, broadly barred; *edge* of *wing shows three notches in flight*; below white, *thickly barred with reddish-brown* and finely lined with blackish; heavy *flight*; notes, high pitched (one resembles that of the Wood Pewee). **Red-shouldered and and Red-tailed Hawks.**

352. BALD EAGLE. 33 in.

(Haliaetus leucocephalus.)

Dark brown, *tail, head and neck white*; yellow bill; edge of wing black; (immature resembles Golden Eagle; adult plumage appears the third year); usually near water; eats fish, carrion, and at times water-birds and mammals; call-notes harsh. **Fish Hawk. Golden Eagle.**

360. AMERICAN SPARROW HAWK. 10½ in.

(Falco sparverius.)

Reddish-brown, usually barred or spotted with black; head bluish, *marked on sides with black*; tail white-tipped with one broad black bar; below white, more or less spotted; narrow pointed wings. FEMALE: much more barred, darker and *streaked* below. *The smallest hawk.*

364. AMERICAN OSPREY. 23 in.

(Pandion haliaetus carolinensis.)

Brown-black, feathers slightly tipped with whitish; *head mottled* with white; *white below*, sometimes faintly spotted; in flight shows dark above, head and lower parts white; nests in colonies along coast; fish-eating; dives for prey; call, a high wailing whistle. **Bald Eagle.**

336. AMERICAN LONG-EARED OWL. 14½ in.

(Asio wilsonianus.)

Very dark brown, lightly mottled with yellowish and white; wings and tail barred; below, streaked on breast, thickly mottled and barred elsewhere; *long, blackish ear-tufts* with light edges; yellow eyes; barking, whirring call; wooded swamp land; beneficial. **Short-eared Owl. Barred Owl.**

Digitized by Google

367. SHORT-EARED OWL. 14½—15½ in.

(Asio flammeus.)

Streaked and mottled dark brown with yellowish; tail broadly barred, also edge of wing; yellowish-white below streaked heavily across breast but *without bars*; *ear-tufts very short*; eyes yellow; grassy marshes and rank undergrowth; sometimes feeds by day in cloudy weather; call shrill; beneficial. **Long-eared Owl. Barred Owl.**

368. BARRED OWL. 18—20 in.

(Strix varia.)

Conspicuously and regularly barred (more finely on head) brown and white with slight yellowish tinge; below white *barred* across breast, *streaked broadly* on belly; *no ear-tufts*; bill *yellow*; eyes almost *black*; woodland; call, a resonant hoot; beneficial. **Short-eared Owl.**

372. SAW-WHET OWL. 7½—8 in.

(Glaux acadica.)

Dark chocolate brown, finely lined on head, spotted on back; tail slightly barred *with white*; below white, thickly streaked; eyes *yellow*; bill black; *no ear-tufts*; the smallest owl common to northeastern America; call resembles "filing a saw"; flies like a woodcock; thickly wooded areas; beneficial. **Screech Owl.**

373. SCREECH OWL. 8—10 in.

(Otus asio.)

Gray to reddish-brown, streaked with black and somewhat mottled; below white, streaked and irregularly *barred*; *ear-tufts*; call, *eerie, wailing tremolo*; frequents orchards; beneficial. **Saw-whet Owl.**

375. GREAT-HORNED OWL. 22 in.

(Bubo virginianus.)

Mottled buff, black and yellow; *long ear-tufts*; eyes *yellow*; below finely barred, *patch on throat white*; wild forested country; very loud, varied calls, "whoo", a scream or bark; attacks poultry and smaller birds but also destroys many mice and rabbits.

Digitized by Google

387. YELLOW-BILLED CUCKOO. 12¼ in.

(Coccyzus americanus.)

Bronzy olive-gray; white below; *lower half of curved bill yellow* tipped with black; outer tail-feathers black tipped *broadly* with white; wings show a *reddish-brown tinge in flight*; glides stealthily from tree to tree and keeps concealed among the leaves; slowly droops and raises tail when perching; song, loud "kuk-kuk, kuk-kuk". **Black-billed Cuckoo. Mourning Dove.**

388. BLACK-BILLED CUCKOO. 12 in.

(Coccyzus erythrophthalmus.)

Bronzy olive-gray; white below; bill *entirely* black; *only a little white* on tail; movements like those of 387; song, soft "coo-coo, coo-coo". **Yellow-billed Cuckoo. Mourning Dove.**

390. BELTED KINGFISHER. 14 in.

(Ceryle alcyon.)

Blue-gray; white collar; *big, crested head*; sits erect watching for fish and flies near the surface of the water, uttering a *rattling* call.

393. HAIRY WOODPECKER. 9 in.

(Dryobates villosus.)

White band down the middle of black back; wings black spotted with white; beneath white; outer tail-feathers white *not barred*; red band on neck of male; large bill; note, sharp resonant "plick". **Downy Woodpecker. Yellow-bellied Sapsucker.**

394c. DOWNY WOODPECKER. 6¼ in.

(Dryobates pubescens medianus.)

Bill small; closely resembles the *Hairy Woodpecker* but is *scarcely larger* than the *English Sparrow* and has outer tail-feathers *barred with black*. **Hairy Woodpecker. Yellow-bellied Sapsucker.**

Digitized by Google

402. YELLOW-BELLIED SAPSUCKER. 8¼ in.

(Sphyrapicus varius.)

Mixed black and white; *yellowish-white underneath*, streaked on sides; white bar *lengthwise* the wing; scarlet patch, bordered with black on crown *and throat* (throat of female white); harsh, squealing note. **Downy Woodpecker. Hairy Woodpecker.**

406. RED-HEADED WOODPECKER. 9½ in.

(Melanerpes erythrocephalus.)

Black and white, not streaked; *entire head, neck and upper part of breast crimson*; pure white belly; note, a loud "kr-r-r-r-r". **Red-bellied Woodpecker.**

409. RED-BELLIED WOODPECKER. 9½ in.

(Centurus carolinus.)

Black and white, back, wings and tail *barred transversely* and very evenly; below grayish-white, *tinged with red*; top of head and back of neck crimson. FEMALE: crown gray. **Red-headed Woodpecker. Hairy Woodpecker.**

412a. NORTHERN FLICKER. 12½ in.

(Colaptes auratus luteus.)

Brown, barred with black; head grayish; back of neck marked with bright red; black crescent on cinnamon-brown spotted breast; in flight shows *white rump* and yellow lining of wings and tail; feeds much *on ground*, unlike other woodpeckers; sharp, nasal call-note; mating call, "flee-ker", repeated. **Meadowlark.**

417. WHIP-POOR-WILL. 9¾ in.

(Antrostomus vociferus.)

Mottled, strongly resembling the bark of certain trees; narrow white band across upper part of breast; big head; in daytime lies *motionless, lengthwise a branch*. **Nighthawk.**

420. NIGHTHAWK. 9½ in.

(Chordeiles virginianus.)

On the wing overhead after insects just before or at dusk; long, pointed wings *show a white patch*; flight resembles that of *Chimney Swift*; sharp note, "skeep"; often nests on the flat roofs of city buildings. **Whip-poor-will.**

Digitized by Google

423. CHIMNEY SWIFT. 5¼ in.

(Chætura pelagica.)

Dark sooty-gray; *bob-tail*; long, pointed wings which often vibrate rapidly in flight; twittering note; never seen perching like other birds. **Swallows.**

428. RUBY-THROATED HUMMING-BIRD. 3¼ in.

(Trochilus colubris.)

Iridescent green and purple; throat of male ruby-red; frequents flowering shrubs and plants; darts here and there and poises in mid air with wings vibrating so rapidly that they make a humming noise; "squeaky note".

444. KINGBIRD. 8½ in.

(Tyrannus tyrannus.)

Dusky black; below white; *tail tipped with broad band of white*; fire-red patch on crown, usually concealed; spreads tail when about to alight; when flying high, may be mistaken for a **Tree Swallow.**

452. CRESTED FLYCATCHER. 9 in.

(Myiarchus crinitus.)

Olive, showing rusty brown on wings and tail; *crested head*; throat and breast, ashy-gray; wing-bars; belly sulphur-yellow; usually perches high; note, hoarse whistle. **Blue Jay.**

456. PHŒBE. 7 in.

(Sayornis phœbe.)

Dull olive-brown, darker on head and tail; whitish below; *bill* and feet black; tail drooping but jerking constantly; note, "phœ-be." **Wood Pewee. Least Flycatcher.**

459. OLIVE-SIDED FLYCATCHER. 7½ in.

(Nuttallornis borealis.)

Olive-gray, wings and tail darker; throat and a narrow line along middle of belly *white*; *large*, *flattened* head; bill *heavy*, black above, light below; *long wings*; tail *notched*; whitish tufts on flanks; bold, conspicuous; plaintive call or loud whistle. **Crested Flycatcher.**

Digitized by Google

461. WOOD PEWEE. 6¼ in.

(Myiochanes virens.)

Dark olive-brown, brownish-black on wings and tail; frowzy head; below white, olive tinge on sides; two narrow white wing-bars; plaintive note repeated slowly and often, "pee-to-way, pee-to-way, hee-rue." **Phœbe. Least Flycatcher.**

463. YELLOW-BELLIED FLYCATCHER. 5½ in.

(Empidonax flaviventris.)

Bright olive-green, tail and wings dark brown; yellowish wing-bars; sulphur-yellow below, *brightest between legs*; tinged elsewhere with pale olive; yellow eye-ring; call, a soft, slow whistle, "pee-a." **Yellow-throated Vireo. Least Flycatcher. Pine Warbler.**

465. GREEN-CRESTED FLYCATCHER. 5¾ in.

(Empidonax virescens.)

Olive-green, wings and tail dark; two *showy* wing-bars; head *broad*; below white, pure on throat and middle of belly, elsewhere palely tinged with greenish-yellow; (immature brighter); open woods near water; common notes, "cheep" or a two-syllabled squeak resembling that made by a rubber doll. **Traill, Least and Yellow-bellied Flycatchers.**

466. TRAILL FLYCATCHER. 5½—6 in.

(Empidonax traillii.)

Olive-green, *tinged with brown*; whitish below, sides grayish; belly slightly yellow; throat *white*; dull, *narrow* wing-bars; pale *eye-ring*; bill, black above, light below; prefers alder growth; a short, soft note, also a longer, sharp, rasping call, accented on second syllable. **Acadian Flycatcher. Least Flycatcher.**

467. LEAST FLYCATCHER. 5¼ in.

(Empidonax minimus.)

Olive-gray, wings, tail and head darker; two narrow wing-bars; *grayish below*, whiter on throat; *jerks its tail* and utters a sharp, energetic call, "che-bec." **Wood Pewee. Phoebe.**

Digitized by Google

474. HORNED LARK. 7¾ in.

(Otocoris alpestris.)

Streaked grayish-brown, somewhat *tinged* with *chocolate*; line *over eye* and *throat yellow* (paler in summer); *breast-patch* and line from bill to and *below eye black*; feathers on sides of crown long, *black*, erectile; tail black, *white-edged*; ground feeder; *runs*; whistling call; in flocks. **Vesper Sparrow. Pipit. Prairie Horned Lark.**

476b. PRAIRIE HORNED LARK. 7¼ in.

(Otocoris alpestris praticola.)

Similar to 474 but smaller; yellow markings almost white; formerly confined to prairies, now spreading eastward. **Horned Lark. Vesper Sparrow. Pipit.**

477. BLUE-JAY. 11½ in.

(Cyanocitta cristata.)

Narrow black collar; *crested*; wings and tail deep blue, white-tipped and black-barred; bold; harsh, noisy notes. **Loggerhead Shrike. Kingfisher. Bluebird.**

488. AMERICAN CROW. 19 in.

(Corvus brachyrhynchos.)

Entirely black, including bill and feet; often seen in flocks; wings appear *frayed* and *ragged in flight*; note, a lusty "caw." **Bronzed Grackle.**

490. FISH CROW. 16 in.

(Corvus ossifragus.)

Most easily distinguished from the common crow by smaller size and *very nasal note*, "car-car-car;" found on the Atlantic coast northward to Connecticut. **Crow.**

494. BOBOLINK. 7½ in.

(Dolichonyx orizivorus.)

Black with light back, creamy buff on neck; long merry musical song; open fields; sings on the wing; FEMALE: sparrow-colored, streaked above.

Digitized by Google

495. COWBIRD. 8 in.

(Molothrus ater.)

Black, head and neck glossy brown; *sparrow-like bill*; fly in flocks, uttering a greasy, squeaking note. FEMALE: brownish. **Bronzed Grackle. Rusty Blackbird.**

497. YELLOW-HEADED BLACKBIRD. 10¼ in.

(Xanthocephalus xanthocephalus.)

Black; head, throat and breast *yellow*; white wing patch. FEMALE: dark brown, yellow markings duller; no wing patch; smaller than male.

498. RED-WINGED BLACKBIRD. 9¼ in.

(Agelaius phœniceus.)

Black showing scarlet shoulder-straps edged with yellow. Likes marshy places; notes "quonk-a-ree," metallic whistle and chattering. FEMALE: much streaked, marked with rusty. **Cowbird. Rusty Blackbird.**

501. MEADOWLARK. 10½ in.

(Sturnella magna.)

Streaked; black crescent on a *bright yellow* breast; *outer tail feathers white*; flies straight, *hovering* as it reaches the ground; sweet, plaintive song; nasal call-note. **Flicker. Dickcissel.**

506. ORCHARD ORIOLE. 7 in.

(Icterus spurius.)

Rich *chestnut-brown*, head, wings and tail black; narrow white wing-bar; high trees; loud, clear song. FEMALE: olivaceous above, greenish-yellow below with two white wing-bars; young male like female but with black throat. **Baltimore Oriole.**

507. BALTIMORE ORIOLE. 7½ in..

(Icterus galbula.)

Bright orange, head, throat, back and wings black; two white wing-bars; loud call, "co-weet, co-weet, co-weet;" song noteworthy. FEMALE: smaller, brownish-yellow showing little black. **Orchard Oriole. Scarlet Tanager.**

Digitized by Google

509. RUSTY BLACKBIRD. 9½ in.

(Euphagus carolinus.)

Black, uniformly iridescent and more or less rusty; tail nearly even. FEMALE: smaller and rustier with pale line over eye. **Bronzed Grackle. Red-wing Blackbird.**

511. PURPLE GRACKLE. 13 in.

(Quiscalus quiscula.)

Back purple to greenish otherwise similar to 511b except that feathers *on back, belly and rump show iridescent bars.* **Bronzed Grackle. Rusty Grackle. Crow.**

511b. BRONZED GRACKLE. 13 in.

(Quiscalus quiscula æneus.)

Iridescent black; head steely blue-green or purple; *body* distinctly bronzy; tail carried rudderwise in flight; pompous walk; rusty, grating call. **Crow.**

517. PURPLE FINCH. 6¼ in.

(Carpodacus purpureus.)

Not "purple," but faded crimson-red, somewhat streaky; belly whitish; *large, heavy bill*; noteworthy song. FEMALE: mottled olive-brown, *resembling sparrows.*

521. AMERICAN CROSSBILL. 6 in.

(Loxia curvirostra minor.)

Male brickish to bright-red, wings and tail dark brown; *bill crossed* and *curved at tip*; feeds on seeds of cone-bearing trees; movements *sluggish* and *parrot-like*; flight, undulating; go in small flocks, singing on the wing. FEMALE: dull olive-green; yellowish rump; belly whitish. **White-winged Crossbill.**

528. REDPOLL. 5½ in.

(Acanthis linaria.)

Streaked dark brown and white; belly, also bars and edges of wing white; sides streaked; *rump*, crown, *throat*, *breast* and sides suffused with rosy-red; blackish throat-patch; bill sharply pointed; canary-like song; in flocks about weedy patches. FEMALE: shows red only on crown; no throat-patch. **Purple Finch. Pine Siskin. Tree Sparrows.**

Digitized by Google

529. AMERICAN GOLDFINCH. 5¼ in.

(Astragalinus tristis).

Canary-yellow, crown, tail and wings black; white wing-bars; canary-like song; also sings in its undulating flight, "per-*chic*-o-ree, per-*chic*-o-ree"; in flocks. FEMALE: olive-tinged, without pure black or yellow. **Yellow Warbler.**

533. PINE SISKIN. 5 in.

(Spinus pinus.)

Black and *buffy-olive, much streaked*; below whitish, streaked; *tail notched*; *slender, pointed bill*; wing and tail-feathers dark-brownish showing *yellow* edges; whitish wing-bars; prefers evergreen or in spring, budding trees; pleasing song. **Redpoll. Female Goldfinch.**

534. SNOWFLAKE. 7 in.

(Plectrophenax nivalis.)

SUMMER PLUMAGE: male, *white*, back, part of wing and inner tail-feathers *black*; female, entirely streaked with black above, showing rusty tinge on wings. WINTER PLUMAGE: similar but male streaked with black on back, showing rusty brown tinge; a ground-feeder; *walker*; goes in flocks; song, sweet and rapid. **Horned Lark. Lapland Longspur.**

540. VESPER SPARROW. 6¼ in.

(Poœcetes gramineus.)

Streaked above and below; underparts whitish, tinged with buff across breast and on sides; shows *two conspicuous white tail feathers in flight*; noteworthy song. **Female Purple Finch.**

542a. SAVANNA SPARROW. 5¼ in.

(Passerculus sandwichensis savanna.)

Much streaked above and below; line over eye and edge of wing, yellowish; cheek sometimes suffused with yellow tinge; *tail short*, feathers pointed; movements stealthy; song, a weak trill. **Henslow Sparrow.**

Digitized by Google

546. GRASSHOPPER SPARROW. 5¼ in.

(Coturniculus savannarum australis.)

Crown black, divided and bordered by buff lines; back broadly, neck finely streaked, black, buff and gray; *tail even, feathers pointed*; below *bright buff*; belly whitish; before eye orange-buff; edge of wing yellow; weak, grasshopper-like song, "chir-r-r-r"; old fences about dry pastures; skulking, mouse-like; flies low and not far. **Savanna and Henslow Sparrows.**

547. HENSLOW SPARROW. 5 in.

(Cotvrniculus henslowii.)

Back conspicuously marked and streaked, showing much chestnut; head and neck a peculiar greenish tinge, *finely* lined with black; below tinged with buff and finely streaked; dark stripes and spots on sides of head; *sharp tail*; hides in the grass and can be carefully observed only with patience. **Grasshopper Sparrow.**

554. WHITE-CROWNED SPARROW. 7 in.

(Zonotrichia leucophrys.)

Ashy-gray; rump, tail and wings brown streaked with black; *crown conspicuously striped* with black and white; no yellow; two white wing-bars; in flocks, often with *White-throated Sparrows*; scratches with both feet; song notable. **White-throated Sparrow.**

558. WHITE-THROATED SPARROW. 7 in.

(Zonotrichia albicollis.)

White throat and wing-bars; unmarked ashy breast; whitish on belly; crown striped with black and white; broad white stripe over the eye shading to yellow in front; yellow and white markings obscure; in flocks about shrubbery; clear, sweet whistle, "pee-bod-dee-dee-dee." IMMATURE: **White-crowned Sparrow.**

559. TREE SPARROW. 6¼ in.

(Spizella monticola.)

Streaked, showing much grayish; two *showy white wing-bars*; chestnut-brown cap; breast whitish, *shading to a dark spot in the middle*; in flocks, often with *Juncos*; a winter visitor; pleasing song. **Chipping Sparrow. Field Sparrow.**

Digitized by Google

560. CHIPPING SPARROW. 5¾ in.

(Spizella passerina.)

Streaked; *plain ashy breast*; two narrow white wing-bars; *chestnut-brown cap*; whitish line over eye; song, a strong clear trill. **Field Sparrow. Tree Sparrow.**

563. FIELD SPARROW. 5¾ in.

(Spizella pusilla.)

Streaked; crown and *bill* reddish-brown; neck ashy; brownish-white *unspotted* breast; two indistinct white wing-bars; song, "fe-u, fe-u, fu, fee, fee, fee," beginning slowly and ending almost in a trill. **Chipping Sparrow. Tree Sparrow.**

567 SLATE-COLORED JUNCO 6¼ in.

(Junco hyemalis.)

Slate color; blackish bib (female brownish) over a white belly; *ivory bill; two white tail feathers*; feeds in flocks on ground, often in company with other birds; "sucks its teeth" for a call-note; song, a melodious trill.

581. SONG SPARROW. 6½ in.

(Melospiza melodia.)

Reddish-brown, showing black streaks; *ashy* line over eye; whitish below with dark brown streaks which form a *spot in the middle of the breast*; cheery song, varied. **Swamp Sparrow. Lincoln Sparrow.**

583. LINCOLN SPARROW. 5¾ in.

(Melospiza lincolni.)

Finely streaked with black and brown; white beneath, with *broad buff band* across the breast and down the sides; narrow streaks on throat, upper breast and sides; dark stripe on each side of throat from corner of mouth; very shy. **Song and Swamp Sparrows.**

584. SWAMP SPARROW. 5¾ in.

(Melospiza georgiana.)

Very darkly colored and heavily streaked; chestnut cap, sometimes indistinctly streaked; *black forehead*; plain ashy breast; shy; frequents bushes along the water's edge. **Song Sparrow. Lincoln Sparrow.**

Digitized by Google

585. FOX SPARROW. 7½ in.

(Passarella iliaca.)

Fox-red back and tail; ashy about neck and head; white breast *thickly* streaked with dark spots; scratches like a hen; alert; noteworthy song. **Brown Thrasher. Hermit Thrush.**

587. TOWHEE. 8¾ in.

(Pipilo erythrophthalmus.)

Black above and black bib (female rich brown); reddish-brown sides; white underneath; *outer tail-feathers white*; scratches about under bushes; brilliant notes, "tow-hee" and "che-wink." **Robin.**

593. CARDINAL. 8½ in.

(Cardinalis cardinalis.)

Bright red, including large blunt bill; *crested*; black markings on face and throat; fine songster. FEMALE: olive-gray, tinged lightly with red and obscurely marked. **Tanagers.**

595. ROSE-BREASTED GROSBEAK. 8¼ in.

(Zamelodia ludoviciana.)

Black, including throat; white rump and wing-patches; *breast white with a brilliant rose-red blotch*; lining of wings also rose-red; *large beak*; sluggish; song, a rich carol. FEMALE: yellowish-brown darkly mottled; no red.

598. INDIGO BUNTING. 5¾ in.

(Passerina cyanea.)

Indigo blue, wings and tail blackish; in flocks; often feeds in the grass; noteworthy song. FEMALE: brown. **Bluebird.**

604. DICKCISSEL. 6 in.

(Spiza americana.)

Streaked; ashy on head and neck; white chin; *black throat-patch*; *yellow breast*; white belly; *line over eye*, *yellow*; wings show chestnut-brown. FEMALE: less yellow; no throat-patch. **Yellow-breasted Chat. Meadowlark.**

Digitized by Google

608. SCARLET TANAGER. 7¼ in.

(Piranga erythromelas.)

Scarlet; *black wings and tail*; rather sluggish; note, "chip, chree;" song resembles that of robin. FEMALE: olive-green; usually near by. **Cardinal. Summer Tanager.**

610. SUMMER TANAGER. 7½ in.

(Piranga rubra.)

Vermilion all over; *no black*. FEMALE: yellowish-olive; usually near the male. **Cardinal.**

611. PURPLE MARTIN. 7½ in.

(Progne subis.)

Blue-black all over; often nests in bird houses; twittering song. FEMALE: lighter, streaked with grayish-white below. **Tree Swallow. Chimney Swift.**

612. CLIFF SWALLOW. 6 in.

(Petrochelidon lunifrons.)

Steel-blue back and crown; *rump*, throat and sides of head *chestnut*; *forehead whitish*; belly white, breast brown-gray marked by steel-blue patch; *tail square*; in flocks; nests in eaves of buildings or under cliffs; twittering call. **Barn and Tree Swallows.**

613. BARN SWALLOW. 7 in.

(Hirundo erythrogaster.)

Steel-blue; chestnut shading to white below; forehead chestnut; *deeply forked tail*. IMMATURE: tail shorter; colors duller. **Cliff Swallow.**

614. TREE SWALLOW. 6¼ in.

(Iridoprocne bicolor.)

Metallic blue-green; *shining white below*; tail appears nearly square in flight; often seen skimming near the surface of still water. **Barn Swallow. Purple Martin.**

Digitized by Google

616. BANK SWALLOW. 5¼ in.

(Riparia riparia.)

White throat and *belly*; *brownish band across breast*, otherwise dull brown-gray; tail notched; nests in holes near top of soft banks (usually sand); twittering call; in flocks. **Rough-winged Swallow. Swift.**

619. CEDAR WAXWING. 7 in.

(Bombycilla cedrorum.)

Smooth snuff-brown; *crested*; chin and line from bill to crest, black; end of tail banded with yellow; in small flocks; "monotonous lisping note."

621. NORTHERN SHRIKE. 10½ in.

(Lanius borealis.)

Gray, very light above eyes and on forehead; below hoary; breast, throat and sides show dusky, *wavy*, *cross-wise bars*; line from bill to eye, continued broadly *below eye* back to nape, *black;* wings and tail black, mostly white-tipped, former marked by white patch; *hooked bill*; straight even flight; seeks commanding perch; song, strong, varied, sometimes mimetic. **Loggerhead Shrike. Blue Jay.**

622. LOGGERHEAD SHRIKE. 9 in.

(Lanius ludovicianus.)

Slaty-gray; commonly white beneath, *unmarked*; wings and tail black, marked with white; *hooked* bill; bar through eye and *between forehead and bill*, black; imitates notes of other birds. **Blue Jay. Catbird.**

624. RED-EYED VIREO. 6 in.

(Vireosylva olivaceus.)

Olive-green; crown slaty; white below, lightly tinged on sides; *dark line above a white one over eye*; clear, persistent song. **Warbling Vireo.**

Digitized by Google

627. WARBLING VIREO. 5¼ in.

(Vireosylva gilva.)

Mouse-gray; olive rump; white below tinged with greenish yellow; creamy-white around eye; song, rippling warble. **Red-eyed Vireo. Least Flycatcher.**

628. YELLOW-THROATED VIREO. 5½ in.

(Lanivireo flavifrons.)

Bright olive-green shading to ashy-blue on rump; *yellow throat* and breast sharply defined against white belly; line around eye and to bill, yellow; *two white wing-bars*; slow, distinct song. **Yellow-bellied Flycatcher. Blue-headed Vireo.**

629. BLUE-HEADED VIREO. 5½ in.

(Lanivireo solitarius.)

Olive-green; *ashy-blue head*; pure white below; sides yellowish; line around eye and to bill, white; *two white wing-bars*; song, noteworthy. **Yellow-throated Vireo.**

631. WHITE-EYED VIREO. 5½ in.

(Vireo griseus.)

Rich olive-green; *two broad yellowish-white wing-bars*; below white, *sides* and *lower belly bright yellow*; white iris; *line from bill to and around eye* yellow; thickets and roadside undergrowth; emphatic call, "whip-tee-you;" mimics notes of other birds. **Yellow-throated and Blue-headed Vireos.**

636. BLACK AND WHITE CREEPER. 5¼ in.

(Mniotilta varia.)

Black and white, *striped*; white belly (female less streaked below); creeps up and down trees like the *Nuthatches*; song rapid, rasping, "he-ho, he-ho, he-ho." **Blackpoll Warbler.**

637. PROTHONOTARY WARBLER. 5¼ in.

(Prothonotaria citrea.)

Entire head, neck and breast, *bright orange-yellow*; blue-gray wings, rump and tail; prefers willows near water; strong, ringing song. FEMALE: duller. **Blackburnian Warbler. Summer Warbler.**

Digitized by Google

641. BLUE-WINGED WARBLER. 4¾ in.
(Helminthophila pinus.)

Bright olive-green; tail and wings ashy-blue, the latter with two *white* bars; *top* of *head*, *forehead* and *under parts bright yellow*, *unmarked*; white tail-spots; *narrow black line through* eye; edge of woods and second growth; two characteristic songs. **Golden-winged and Brewster Warblers.**

642. GOLDEN-WINGED WARBLER. 5 in.
(Helminthophila chrysoptera.)

Blue-gray; crown and *patch on wing*, golden yellow; *throat* and stripe through eye, *black*; white below; song, drawling "zee, zee, zee, zee." **Blue-winged and Parula Warblers.**

645. NASHVILLE WARBLER. 4¾ in.
(Helminthophila rubricapilla.)

Olive-green; head and neck bluish gray; indistinct brown patch on crown; *bright yellow below*; no wing-bars; *white eye-ring*; song, not loud, "wee-see, wee see, wit-a-wit." **Tennessee Warbler.**

647. TENNESSEE WARBLER. 4½ in.
(Helminthophila peregrina.)

Olive-green, brightest on rump; head and neck, blue-gray; *dull white below*; faint white line over eye; no wing-bars; song, shrill and feeble. **Nashville Warbler.**

648a. NORTHERN PARULA WARBLER. 4½ in.
(Compsothlypis americana usneae.)

Grayish-blue, *yellowish across the back*; throat and breast yellow, latter crossed by brownish band; wing-bars; peculiar song. **Golden-winged Warbler.**

650. CAPE MAY WARBLER. 5¼ in.
(Dendroica tigrina.)

Yellowish-olive spotted with black on back; crown very dark; cheeks yellow with an *orange-brown ear-patch*; yellow below, black streaks on breast; wing-bars; song all on one note. **Black-throated Green Warbler. Magnolia Warbler.**

Digitized by Google

652. YELLOW WARBLER. 5 in.

(Dendroica æstiva.)

Yellow all over shading to olive on back; wings and tail darker, showing yellow edgings; orange-brown streaks on breast; song, penetrating and unceasing; nests often in city trees. FEMALE: not so yellow, less streaked. **American Goldfinch.**

654. BLACK-THROATED BLUE WARBLER. 5¼ in.

(Dendroica cærulescens.)

Slaty-blue; below white; black throat and sides; *white patch on wing*; *hoarse* note, "kree, kree." FEMALE: grayish; *no black* but may be recognized by *wing patch*. **Blue-gray Gnatcatcher.**

655. MYRTLE WARBLER. 5½ in.

(Dendroica coronata.)

Blue-gray, streaked; crown, *rump* and spot on each side of breast *yellow*; white below, marked on breast and sides with black; usually in small flocks. **Magnolia Warbler.**

657. MAGNOLIA WARBLER. 5 in.

(Dendroica magnolia.)

Back and tail black; crown blue-gray; yellow rump; *yellow* below, streaked with black on breast and sides; white patch on wings; broad white band across tail, *seen from below*. **Myrtle Warbler. Cape May Warbler.**

658. CERULEAN WARBLER. 4½ in.

(Dendroica cærulea.)

Azure-blue, marked with black on back and sides of head; white below, streaked with blue; two white wing-bars; frequents tree-tops; song resembles that of *Parula*. **Blue-gray Gnatcatcher.**

659. CHESTNUT-SIDED WARBLER. 5 in.

(Dendroica pensylvanica.)

Back shows black streaks mixed with olive-green; yellow cap; white below; *chestnut markings* on *sides of breast*; wing-bars; song, chee, chee, chee, *chee*-ar." **Bay-Breasted Warbler.**

Digitized by Google

660. BAY-BREASTED WARBLER. 5½ in.

(Dendroica castanea.)

Ashy-brown, black-streaked; black cheeks and forehead; throat, upper part of breast, sides and crown, *rich chestnut-brown*; white wing-bars; song, soft, liquid warble. **Chestnut-sided Warbler.**

661. BLACK-POLL WARBLER. 5½ in.

(Dendroica striata.)

Streaked black and white; *black crown*; white cheeks; below white with line of black spots down sides; tree-tops; song, a staccato musical "chink" repeated rapidly. **Black and White Creeper.**

662. BLACKBURNIAN WARBLER. 5¼ in.

(Dendroica blackburniæ.)

Black, streaked with white; middle of crown, throat and sides of head *flaming orange*; black patch behind eye; white wing-bars; usually, high up in evergreen trees; shrill song; FEMALE: duller. **Prothonotary Warbler.**

667. BLACK-THROATED GREEN WARBLER. 5¼ in.

(Dendroica virens.)

Olive-green; *bright yellow cheeks*; *throat and upper part of breast black*; *tail shows much white*; white wing-bars; song, repeated often and rapidly, "chee, chee, chee, char, chee." **Cape May Warbler. Hooded Warbler.**

671. PINE WARBLER. 5¾ in.

(Dendroica vigorsii.)

Olive-green; yellow breast; sides of head suffused with yellowish, sometimes forming a *yellow line over eye*; two dull white wing-bars; song, a loud, penetrating trill. **Yellow-bellied Flycatcher. Female Maryland Yellow-throat. Female Wilson Black-cap.**

Digitized by Google

672. PALM WARBLER. 5 in.

(Dendroica palmarum.)

Streaky olive-gray; *chestnut cap*; yellow line over eye; beneath yellowish (brightest on throat and under tail) streaked with brown; feeds on ground in flocks; *tilts its tail* constantly. **Water-Thrushes. Yellow Palm Warbler.**

672a. YELLOW PALM WARBLER. 5½ in.

(Dendroica palmarum hypochrysea.)

Similar to 672 but *uniform* yellow beneath and more strongly olive above; song, soft, broken trill, "tzee-tzee". **Palm Warbler.**

673. PRAIRIE WARBLER. 4¾ in.

(Dendroica discolor.)

Olive-green, *middle of back spotted with chestnut*; sides of head, broad line over eye and under parts, bright yellow; *lines through and below eye* with *chain of streaks down sides black*; white *tail-spots* and yellowish wing-bars; in wild pastures; song, thin, "zee", repeated on ascending scale. **Magnolia, Canadian, Cape May and Female Black-throated Green Warblers.**

674. OVEN-BIRD. 6¼ in.

(Seiurus aurocapillus.)

Olive-green showing yellow tinge; below pure white, thickly spotted; *orange-brown cap* bordered with black stripes; scratches like a hen; tilts tail; under shrubbery; note, "teacher" repeated rapidly with increasing emphasis. **Water-Thrushes. Wood Thrush.**

675. WATER-THRUSH. 6 in.

(Seiurus noveboracensis.)

Plain olive-brown; below white, tinged with sulphur-yellow and thickly streaked, finely *on chin* and *throat*; belly plain; *short bill, yellowish* line over eye; *wags tail*; noticeable song, high, ending low. **Oven-bird. Other Water-Thrushes.**

Digitized by Google

675a. GRINNELL WATER-THRUSH. 5¾ in.

(Seiurus noveboracensis notabilis.)

Very dark olive-brown; below white, palely tinged with yellowish, thickly streaked; *pale* stripe over the eye; skulks along the very edge of the water; *tilts tail*; brilliant song. **Louisiana Water-Thrush. Oven-bird. Palm Warbler.**

676. LOUISIANA WATER-THRUSH. 6¼ in.

(Seiurus motacilla.)

Olive; white below, buff-tinged and streaked with black; *throat and middle of belly unmarked*; *white line over eye*; loud, ringing song. **Grinnell Water-Thrush. Oven-bird. Palm Warbler.**

678. CONNECTICUT WARBLER. 5½ in.

(Oporornis agilis.)

Olive-green; head ashy, throat darker; *white ring* around eye; belly yellow; wings long and pointed; low bushes and swampy places; penetrating song. **Mourning Warbler.**

679. MOURNING WARBLER. 5½ in.

(Oporornis philadelphia.)

Bright olive-green; head, neck, throat and upper part of breast *bluish-ash*; yellow belly; black band on breast; about bushes and in the grass; noteworthy song. **Connecticut Warbler.**

681. MARYLAND YELLOW-THROAT. 5¼ in.

(Geothlypis trichas.)

Olive-green showing brownish tinge; *broad, black stripe across forehead and through eye*, bordered with grayish-white; yellow beneath, lighter on belly; wings short; in low bushes; song, "witch-i-ty, witch-i-ty, witch-i-ty." FEMALE: duller, *without black stripe* and easily confused with **Female Mourning, Connecticut and Pine Warblers. Kentucky and Wilson Warblers.**

Digitized by Google

683. YELLOW-BREASTED CHAT. 7¼ in.

(Icteria virens.)

Olive-green; bright yellow below, belly *white*; region about eye blackish, marked with a white line above eye; white spot and shorter line below eye; the largest warbler; remarkable song. **Dickcissel. Maryland Yellow-throat. Yellow-Throated Vireo.**

684. HOODED WARBLER. 5½ in.

(Wilsonia citrina.)

Olive-green; yellow below; head, neck and upper breast black; *yellow "mask" across forehead and cheeks*; white spots on tail; FEMALE: less black. **Maryland Yellow-throat. Wilson Warbler. Canadian Warbler.**

685. WILSON WARBLER. 4¾ in.

(Wilsonia pusilla.)

Bright olive-green; yellow below; *black cap*; in low bushes. **Pine Warbler. Female Maryland Yellow-throat.**

686. CANADIAN WARBLER. 5½ in.

(Wilsonia canadensis)

Bluish-lead color; yellow below with *a necklace of black spots*; no wing-bars; note-worthy song. **Magnolia Warbler.**

687. AMERICAN REDSTART. 5¼ in.

(Setophaga ruticilla.)

Male black, with *six flame-colored spots* (female olive, with *six yellow spots*); spreads tail like a fan; song, "chee, chee chee, chew;" darting motions.

697. AMERICAN PIPIT. 6 in.

(Anthus rubescens.)

Streaked grayish-brown; below buff, breast and sides darkly streaked; tip of wing and *outer edge of tail whitish*; light line above and ring around eye; *long hind toe-nail*; *wagging walk*; undulating flight; in flocks; song, soft "dee-dee". **Larks. Water-thrushes.**

Digitized by Google

703. MOCKINGBIRD. 10½ in.

(Mimus polyglottus.)

Ashy-gray, wings and tail darker; whitish below; *wing shows white patch*; outer tail-feathers white; famous for its song. **Catbird. Loggerhead Shrike. Cuckoos.**

704. CATBIRD. 8¾ in.

(Dumatella carolinensis.)

Slate-gray; bill, crown and tail black; *chestnut patch under tail*; about shrubbery; willing to be looked at; calls like a cat; song, greatly varied.

705. BROWN THRASHER. 11¼ in.

(Toxostoma rufum.)

Bright rusty-red; white below, much streaked; long bill and *long, sweeping tail*; wing-bars; song, loud and varied; about bushes. **Wood Thrush.**

718. CAROLINA WREN. 5½ in.

(Thryothorus ludovicianus.)

Reddish-brown, brightest on rump; below rusty-buff; throat and *line over eye*, whitish; wings and tail barred, the latter long and carried erect; "quick as a flash," flight jerky; very loud, ringing song and striking call-notes; near wooded streams in wild pasture undergrowth. **Bewick and Long-billed Marsh Wrens.**

721. HOUSE WREN. 5 in.

(Troglodytes aedon.)

Brown, barred with darker ; below grayish, tinged with brown, also barred; rusty-colored tail, *two inches long*; remarkable song. **Other Wrens.**

722. WINTER WREN. 4¼ in.

(Nannus hiemalis.)

Dark tawny-brown, barred on wings and tail; lighter below, barred except on throat and breast; pale line over eye; *stub tail, carried erect*; fine songster. **Other Wrens.**

Digitized by Google

725. LONG-BILLED MARSH WREN. 5¼ in.

(Telmatodytes palustris.)

Brown, almost black on head, tail and between shoulders, *the latter streaked with white*; barred only on wings and tail; *line over eye* and middle of breast white; *long bill*; spluttering song. **Other Wrens.**

726. BROWN CREEPER. 5½ in.

(Certhia familiaris americana.)

Brown finely streaked with white; silky white below; long curved bill; *flies to the base of tree-trunks and works up in spirals* bracing with its long pointed tail; calls softly, "screep, screep." **Nuthatches.**

727. WHITE-BREASTED NUTHATCH. 5¾ in.

(Sitta carolinensis.)

Ashy-blue; cap and back of the neck, black; white below; constantly exploring the bark of trees; does not brace,woodpecker fashion, with its tail; note, nasal "yank, yank, yank." **Red-breasted Nuthatch. Brown Creeper.**

728. RED-BREASTED NUTHATCH. 4¾ in.

(Sitta canadensis.)

Bluish-gray; crown and *stripe through eye, black* with a white line over eye; *below rusty-red*, throat white; climbs up and down and around trees; note, "cark, cark, cark." **White-breasted Nuthatch.**

731. TUFTED TITMOUSE. 6¼ in.

(Baeolophus bicolor.)

Ashy-gray; below whitish; sides reddish-brown; *head crested*; *bill* and band above *black*; loud emphatic song; whistling call "peto." **Chickadee. Catbird.**

735. CHICKADEE. 5¼ in.

(Penthestes atricapillus.)

Ashy-gray; *black throat* and *cap*; white cheeks; frowzy; lively; often swings head downward from the tips of twigs; call-note "chick-a-dee-dee"; song musical, "pewee," confused with note of *Phœbe*.

Digitized by Google

748. GOLDEN-CROWNED KINGLET. 4 in.

(Regulus satrapa.)

Olivaceous; two whitish wing-bars; *orange-yellow stripe edged with black on crown*; restless: often seen feeding in evergreens; note, rapid "tzee, tzee, tzee." **Ruby-crowned Kinglet.**

749. RUBY-CROWNED KINGLET. 4¼ in.

(Regulus calendula.)

Dark greenish-olive; light underneath; *ring about eye* and wing-bars white; male shows at will a brilliant ruby crown-patch; remarkable song. **Golden-crowned Kinglet. Warblers.**

751. BLUE-GRAY GNATCATCHER. 5 in.

(Polioptila cærulea.)

Blue-gray; white beneath; *long tail* edged with white; male has a black line over eye; fidgety, never long in a place; floating flight; wheezy song. **Black-throated Blue Warbler. Cerulean Warbler.**

755. WOOD THRUSH. 7¾ in.

(Hylocichla mustelina.)

Bright brown on head shading to olive-brown on rump and tail; white beneath with conspicuous *round* black spots except on throat and middle of belly; remarkable song. **Other Thrushes. Fox Sparrow.**

756. WILSON THRUSH. 7¼ in.

(Hylocichla fuscescens.)

Uniform tawny brown; no ring about eye; below white, faintly tinged and lightly spotted; shy, about bushes; notable song. **Other Thrushes.**

757. GRAY-CHEEKED THRUSH. 7½ in.

(Hylocichla aliciae.)

Uniform olive; below white, *very palely* suffused with buff; cheeks *gray tinged*; breast and sides of throat spotted; **Olive-backed and other Thrushes.**

Digitized by Google

758a. OLIVE-BACKED THRUSH. 7¼ in.

(Hylocichla ustulata.)

Uniform dark olive; below white, darker on sides; cheeks, throat, breast and ring about eye buff; spotted on breast; song loud and ringing. **Gray-cheeked Thrush. Hermit Thrush. Wilson Thrush.**

759b. HERMIT THRUSH. 7¼ in.

(Hylocichla guttata pallasii.)

Olive-brown; *tail reddish-brown*; below dull white tinged with buff and thickly spotted across the breast; shy, hiding in bushes; famous for its song but silent during migration. **Other Thrushes.**

761. ROBIN. 10 in.

(Planesticus migratorius.)

Dark slate color; black head; *yellow bill*; throat white streaked with black; underparts bright chestnut-red; sings "cheer-i-ly, cheer-i-ly, cheer-up." IMMATURE: spotted with black. **Towhee.**

766. BLUEBIRD. 6½ in.

(Sialia sialis.)

Sky-blue; brownish-red below; female shows less blue; usually in pairs; call, "pu-ri-ty, pu-ri-ty;" often heard before seen. **Blue Jay.**

Digitized by Google

Note to Key.

Since no key can make up for inaccurate observation, it is essential that at least one or two points about an unknown bird shall have been *clearly seen* before turning to any key for aid in identification.

Keep in mind (1) the *relative size* and (2) see accurately some one or more distinguishing markings of any bird in question.

The following key *includes the land birds* described in the foregoing pages. These are arranged by number, first in three groups according to size (using the Robin and English Sparrow as 10 and 6 inch measures of length respectively) and second, according to distinguishing colors or markings.

Suppose a bird at least as large as a Robin, i. e., 10 inches or more in length, is seen, showing white outer tail-feathers and some yellow beneath. Upon referring to the key only one number, namely 502, will be found which answers this description. When more than one number fulfils the conditions of the observation it becomes necessary to compare descriptions before coming to a decision.

Blackface figures indicate female birds that are noticeably unlike the corresponding males.

Digitized by Google

BRIGHT COLORS. STREAKED, BARRED OR SPOTTED	Showing BLUE .....
	Showing RED .....
	Showing YELLOW .....
	Showing BLACK .....
	Showing BROWN .....
BRIGHT COLORS. *NEVER* STREAKED, BARRED OR SPOTTED	Showing BLUE .....
	Showing RED .....
	Showing YELLOW .....
	Showing BLACK .....
	Showing BROWN .....
IRIDESCENT COLORS	.....
DULL COLORS	FEW MARKINGS
	Sharply defined.....
	Obscure .....
	NO MARKINGS.....
BACK	STREAKED
	Entirely streaked or mottled, no distinct cap....
	Rump plain, cap or striped crown .....
	Gray or ashy about head or neck.....
	BARRED AND SPOTTED
	Barred, including wings.....
	Black and white, wings barred or spotted.....
	NO MARKINGS
	Back and wings alike.....
	DISTINCTIVE MARKINGS
	Rump patch white, yellow or brown .....
	On back of neck, between shoulders or middle of back.....

Digitized by Google

10 inches or more	Between 6 and 10 inches.	Less than 6 inches.
390, 477.....	.....	658.....
412a, 761.....	406, 409, 517, 521, 595.....	428, 528, 687 .....
412a, 501.....	604.....	650, 652, 655, 657, 662, 667, 671
.....	.....	672a, 681 686, 687.....
360, 412a, 477, 501,	406, 409, 493, 494, 534, 604,	533, 650, 655, 657, 659, 660, 662,
.....	.....	667, 686, 687.....
289, 331, 339, 343,	585, 755, 756, 759b.....	598, 659, 660, 718 721, 722, 725
360, 705.....	.....	726.....
.....	766.....	598, 629, 642, 648, 654, 727, 728
.....	.....	751.....
761.....	498, 593, 608, 610, 766.....	.....
497.....	498, 507, 619, 683.....	529, 628, 637, 642, 648a, 671, 678
.....	.....	679, 681, 684, 685.....
488, 490, 497.....	444, 495, 498, 509, 611, 613,	428, 654.....
.....	614.....	.....
.....	495, 506, 587, 619, 766.....	728.....
316, 511, 511b.....	493, 495, 509, 611, 612, 613,	428.....
.....	614.....	.....
316, 703.....	417, 444, 567, 622, 704.....	463, 683, 735, 748, 749.....
364, 387, 388, 497,	456, 459, 461, 495, 509, 593	466, 467, 529, 598, 627, 647, 749..
621.....	608, 611, 624.....	.....
488.....	456, 495, 509, 610, 611.....	423, 598, 627, 681.....
501.....	373, 417, 420, 494, 517, 540,	533, 542a, 546, 547, 583, 636, 650,
.....	581, 595.....	661.....
	558.....	560, 662, 726.....
.....	554, 559, 585, 604.....	563, 584.....
289, 300, 360, 368,	372, 409.....	721, 722 .....
412a.....	.....	.....
.....	393, 394c, 402.....	636.....
316, 387, 388, 488,	456, 495, 567, 611, 624, 674,	423, 627, 645, 647, 675a, 678, 679,
511b, 761.....	676, 683, 756, 757, 758a...	681, 684, 685, 686, 728, 735.....
331, 412a.....	595, 612.....	650, 655, 657, 725.....
412a.....	393, 394c, 402, 409.....	547, 648, 673, 725.....

Digitized by Google

UNDER PARTS	No Markings—Pure white, yellow or ashy
	Dull colors, indistinctly shaded
	Few, if any Markings
	Distinctive red, yellow or brown breast
	Belly white or yellow, sharply contrasting with breast
	Distinctive Markings
	A line of streaks down sides or across breast
	Spotted or finely streaked with throat and middle of belly plain
	Bib, collar or band across breast
	Spot or patch on throat, sides, breast or sides of neck
	Tinged
	Noticeably with yellow, red, buff or brownish
HEAD	Top
	Crested or red crown-patch displayed at will
	Crown conspicuously striped
	Distinct cap, no line over eye
	Distinct cap, with line over, through or back of eye
	Forehead showing black, sometimes with bar through eye
	Hood
	Sides
	Eartufts
	Conspicuous line over or through eye
	Inconspicuous but distinctive line over eye
	Conspicuous eye-ring or, black or yellow mask
	Distinctly marked or, colored on cheeks or sides of head
	Neck band
	Bill.
	Hooked, or noticeably long, sometimes curved
	Large and stout, or noticeably short and thick
	Used for hammering

Digitized by Google

10 inches or more.	Between 6 and 10 inches.	Less than 6 inches.
387, 388,	393, 394a, 406, 444, 461, 463	316, 466, 546, 563, 584, 627, 629,
	554, 614, 624, 683	641, 642, 645, 658, 678, 726, 727,
		751
703	456, 461, **506**, **507**,**509**, 622,	467, 563, 584, 647, 725, 735, 748,
		749, 751
501, 761	613, 683, 766	628, 637, 645, 671, 681, 684, 685,
	452, 567, 587, 604, 612, 683,	628, 631, 654, 678, 679
	402, 585	655, 657, 658, 659, 661, 662, 673
		686
360	417, 420, 567, 587, 655, 676,	547, 583, 648a, 654, 658, 660, 667,
		684
390, 412a	417, 587, 616	583, 667, 679
375, 412a, 501	402, 474, 558, 559, 581, 595,	428, 528, 642, 679, 689, 735
	604, 612	
	402, 409, 452, 517, 675, 757	463, 528, 547, 563, 583, 629, **671**,
	758a	672, 672a, 722, 728
289, 390, 477	402, 444, 452, 593, 619, 731,	749
	331, 554, 558, 674	546, 636, 748
	409, 704	529, 641, 657, 660, 661, 685, 729
		735
	559	560, 563, 642, 655, 659, 672, 728
	593, 619	584, 660, 681, 751
352, 497	406, 495, 506, 507	629, 637, 679, 684
300, 366, 375	373, 474	
289, 501	393, 394c, 474, 474b, 558,	628, 641, 642, **650**, 657, 667, 671,
	604, 622, 675, 676, 683	673, 675a, 718, 725, 751
497	509, 624, 675a	542a, 647, **658**, 672, 697
	683, 758a	463, 749, 681, 628, 645, 629, 684
		678, 631
316, 360, 621	409, 619, 731, 757	547, 583, 642, 650, 658, 660, 661,
		662, 667, 735
390, 477	393, 494	
331, 360, 387, 388,	521, 622	428, 725, 726
621, 705		
	495, 517, 593, 595	529, 598
412a	393, 394c, 402, 406, 409, 412a	

Digitized by Google

WINGS	Showing notches in flight
	Long and pointed
	One or two conspicuous wing-bars
	Spots, patches or bright markings
TAIL	MARKINGS
	Outer feathers conspicuously white or spotted
	Reddish brown or, tipped with white or yellow
	Barred or, banded with black or white
	SIZE
	Very long, sometimes keeled or forked
	Very short
	MOVEMENTS
	Used for bracing
	Jerking or tilting motions
	SHAPE
	Square
	Rounded
	Notched
	Narrow and pointed

Digitized by Google

10 inches or more.	Between 6 and 10 inches.	Less than 6 inches.
339, 343, 488.....	.....	.....
.....	417, 420, 611, 613, 614.....	423.....
705.....	452, 461, 506, 507, 554, 559	463, 465, 467, 528, 529, 628, 629
.....	.....	631, 641, 648a, 658, 659, 660, 661,
.....	.....	667, 671, 673, 726, 749.....
393, 497, 621, 703,	402, 406, 420, 498, 587, 595,	642, 654, 655, 657, 687.....
.....	622.....	.....
387, 390, 501, 703.	393, 394c, 474, 540, 567, 587,	655, 658, 660, 667, 671, 672, 687..
761.....	622.....	697, 751.....
316, 332, 333, 337.	444, 452, 585, 619, 759b....	.....
339, 360, 387, 477	.....	.....
300, 331, 332, 360.	394c, 402, 409.....	657, 721, 722, 725.....
366, 477.....	.....	.....
316, 387, 388, 511b.	613.....	726, 751.....
705.....	.....	.....
.....	412a.....	423, 428, 721, 722, 727, 728.....
412a.....	393, 394c, 402, 406, 409.....	726.....
.....	456, 674, 675, 675a, 676.....	467, 672, 672a, 675a, 697.....
332.....	612, 614.....	546.....
333.....	.....	.....
.....	459, 613, 616.....	533.....
316.....	.....	542a, 547.....

Digitized by Google

Table of Occurrence.

In this table the birds observed by the authors in Lincoln Park, Chicago, are arranged according to the order of their comparative frequency. The abundance of the birds, that is, the number of individual birds of different species, is in no wise shown but simply the relative representation of each species. For instance, out of a total of 454 mornings covering observations during six years, the *Robin* was seen on 383 different mornings. The record for 1902, during the absence of the authors, was made by Mr. H. V. Bozell.

Digitized by Google

		NUMBER OF MORNINGS SEEN IN LINCOLN PARK, CHICAGO						
		1898	1899	1900	1901	1902	1903	Total
1	Robin	62	52	57	69	61	82	383
2	Bronzed Grackle	57	58	57	60	57	72	361
3	Blue Jay	49	58	61	65	40	51	324
4	Cowbird	41	34	44	44	39	54	256
5	Junco	33	35	32	32	51	42	225
6	Towhee	35	18	24	44	45	48	214
7	Song Sparrow	27	33	19	38	37	36	190
8	Flicker	29	18	21	33	37	46	184
9	Ruby-crowned Kinglet	31	28	23	30	22	39	173
10	White-throated Sparrow	30	23	23	31	26	35	168
11	Bluebird	26	1	7	31	35	50	150
12	Wilson Thrush	34	35	23	21	11	20	144
13	Yellow-bellied Sapsucker	23	12	15	30	27	34	141
14	Yellow Warbler	27	30	24	18	13	27	139
15	Belted Kingfisher	25	21	24	27	21	19	137
16	Redheaded Woodpecker	23	23	16	26	16	26	130
17	Myrtle Warbler	19	25	11	21	22	31	129
18	Brown Thrasher	27	8	20	21	16	29	121
19	Brown Creeper	18	15	19	14	28	27	121
20	Catbird	22	27	16	17	10	17	109
21	Golden-crowned Kinglet	18	14	13	26	15	23	109
22	Chipping Sparrow	25	11	11	12	17	32	108
23	American Goldfinch	21	19	12	20	5	18	95
24	Tree Swallow	13	11	16	19	13	22	94
25	American Redstart	18	16	19	8	10	20	91
26	Barn Swallow	15	14	10	19	14	19	91
27	Fox Sparrow	11	9	14	9	27	21	91
28	Chimney Swift	16	11	14	15	11	22	89
29	Blue-gray Gnatcatcher	18	16	11	10	16	17	88
30	White-crowned Sparrow	9	5	18	22	16	17	87
31	Palm Warbler	8	21	7	11	19	20	86
32	Hermit Thrush	13	9	10	18	9	27	86
33	Black and White Creeper	15	15	11	13	12	18	84
34	Least Flycatcher	18	18	8	14	7	18	83
35	Magnolia Warbler	13	13	13	5	16	20	80
36	B'k-throated Blue Warbler	12	14	9	13	12	16	76
37	Olive-backed Thrush	8	4	14	16	11	23	76
38	Phoebe	15	4	5	16	15	19	74
39	Baltimore Oriole	18	6	10	15	11	13	73
40	Grinnell Water-Thrush	17	6	9	14	8	17	71
41	Field Sparrow		2	12	17	23	17	71
42	Swamp Sparrow	4	4	14	12	17	17	68
43	Crow	4	11	7	11	16	18	67

Digitized by Google

		NUMBER OF MORNINGS SEEN IN LINCOLN PARK, CHICAGO						
		1898	1899	1900	1901	1902	1903	Total
44	Chestnut-sided Warbler	11	6	11	13	9	16	66
45	Oven-bird	9	11	11	12	11	11	65
46	White-breasted Nuthatch	12	6	17	21	6	2	64
47	Maryland Yellow-throat	18	17	10	8	3	15	61
48	Wood Pewee	12	14	9	6	7	13	61
49	Downy Woodpecker	14	7	4	8	17	10	60
50	House Wren	1	...	10	11	13	24	59
51	Red-eyed Vireo	13	10	5	6	6	16	56
52	Loggerhead Shrike	5	1	16	13	7	13	55
53	Indigo Bunting	12	17	2	6	5	11	53
54	Bl'k-throated Green Warbler	4	9	4	9	6	12	44
55	Blackburnian Warbler	11	4	9	7	5	6	42
56	Black-poll Warbler	6	7	8	3	3	15	42
57	Kingbird	8	1	5	3	6	17	40
58	Scarlet Tanager	7	3	7	4	3	10	34
59	Ruby-thr. Humming-bird	7	9	4	3	1	10	34
60	Mourning Dove	5	3	6	10	2	8	34
61	Rose-breasted Grosbeak	5	2	6	8	6	6	33
62	Tree Sparrow	...	...	10	3	12	8	33
63	Winter Wren	3	5	11	...	3	10	32
64	Meadowlark	3	...	7	13	5	4	32
65	Purple Martin	1		...	6	4	20	31
66	Cape May Warbler	3	2	9	2	6	7	29
67	Warbling Vireo	7	1	4	3	4	10	29
68	Mourning Warbler	5	8	2	4	4	5	28
69	Wilson Black-cap Warbler	9	1	3	2	5	8	28
70	Yellow-billed Cuckoo	8	7	1	3	1	7	27
71	Crested Flycatcher	5	2	...	1	10	9	27
72	Bobolink	2	...	9	10	1	4	26
73	Canadian Warbler	8	3	6	4	1	4	26
74	Red-winged Blackbird	4	...	2	8	3	7	24
75	Wood Thrush	8	3	...	5	3	3	22
76	Orchard Oriole	4	3	5	1	1	6	20
77	Red-breasted Nuthatch	...	...	10	...	3	7	20
78	Marsh Hawk	...	2	1	8	5	4	20
79	Hairy Woodpecker	3	1	4	...	4	7	19
80	Bay-breasted Warbler	1	1	2	1	4	10	19
81	Parula Warbler	6	...	...	2	9	2	19
82	Chickadee	3	...	7	8	...	...	18
83	Pine Warbler	...	...	3	1	5	9	18
84	Henslow Sparrow	3	2	1	2	3	6	17
85	Nashville Warbler	1	1	4	1	...	9	16
86	Yellow-throated Vireo	2	...	2	2	3	7	16

Digitized by Google

		NUMBER OF MORNINGS SEEN IN LINCOLN PARK, CHICAGO						
		1898	89	1900	1901	1902	1903	Total
87	Cedar Waxwing	6	5	...	1	2	1	15
88	Purple Finch	4	...	4	...	1	5	14
89	Vesper Sparrow	...	...	6	...	4	3	13
90	Blue-headed Vireo	...	...	2	1	6	4	13
91	Gray-cheeked Thrush	...	...	...	...	3	10	13
92	American Sparrow Hawk	1	1	...	4	2	4	12
93	Yellow-bellied Flycatcher	...	...	5	...	2	4	11
94	Nighthawk	...	2	1	1	...	7	11
95	Lincoln Sparrow	...	...	1	...	1	8	10
96	Black-billed Cuckoo	...	1	2	2	2	2	9
97	Rusty Blackbird	...	2	...	3	...	2	7
98	Yellow-breasted Chat	...	...	...	...	1	6	7
99	Long-billed Marsh Wren	1	2	...	...	...	3	6
100	Louisiana Water-Thrush	...	...	1	...	3	2	6
101	Prothonotary Warbler	...	1	...	...	3	1	5
102	Savanna Sparrow	...	...	...	...	...	5	5
103	Blue Golden-winged Warb	1	...	1	1	1	..	4
104	Whip-poor-will	1	1	...	1	...	1	4
105	Connecticut Warbler	...	...	...	...	...	3	3
106	Dickcissel	...	...	...	1	2	...	3
107	Tennessee Warbler	2	...	...	...	...	...	2
108	Mockingbird	...	...	...	2	...	...	2
109	Cerulean Warbler	...	...	1	...	1	...	2
110	Summer Tanager	...	1	...	...	...	...	1
111	Cardinal	...	...	1	...	...	...	1
112	Red-bellied Woodpecker	1	...	...	...	...	...	1
113	Hooded Warbler	...	...	...	...	...	1	1
114	Yellow-headed Blackbird	...	...	...	1	...	...	1
	Total number of days	66	77	74	78	72	87	454
Av'ge No. of kinds seen daily		18.83	12.75	14.59	16.23	16.45	19.53	16.34

Digitized by Google

Table of Arrival.

NOTE:—The following table gives the average date of the first arrival of each bird together with the data from which it was reckoned, for the seven years from 1897 to 1903 inclusive, *as observed in Lincoln Park*, Chicago, by the authors. The record for 1902, in the absence of the authors, was kept by Mr. Harold V. Bozell.

Attention is called to the fact that many birds which are common in the neighboring country are rare in the Park. Among these may be mentioned the *Red-winged Blackbird, Bobolink, Warbling Vireo, Prairie Horned Lark and Tufted Titmouse.*

Digitized by Google

	Average First Appearance	MARCH							APRIL							MAY						
		1897	1898	1899	1909	1901	1902	1903	1897	1898	1899	1900	1901	1902	1903	1807	1898	1899	1900	1901	1902	1903
Family COLUMBIDAE																						
Mourning Dove	April 15.			25								6	10	26	1		9					
Family FALCONIDAE.																						
Marsh Hawk	April 12.					25					25	19		21	1							
Sparrow Hawk	April 17.					24	21				25	18			30		18					
Family CUCULIDAE																						
Yellow-billed Cuckoo	May 14																19	1	16	17	17	12
Black-billed Cuckoo	May 16																	27	12	15	16	12
Family ALCEDINIDAE																						
Belted Kingfisher	April 1.	28	25	22		26						10		11	10							
Family PICIDAE																						
Hairy Woodpecker	April 11.						25	6		11		19						24				
Downy Woodpecker	March 30			27			10	6	7	8		25	5									
Yellow-bellied Sapsucker	April 3		30			19	27		25		8	6			1							
Red-bellied Woodpecker	May 22																22					
Red-headed Woodpecker	April 29								26	29	28		27	27	28				6			
Flicker	March 31		25			23	24	18	15		11	7										
Family CAPRIMULGIDAE																						
Whip-poor-will	May 12											30					21			9		17
Nighthawk	May 15																	10	13	18	18	18
Family MICROPODIDAE																						
Chimney Swift	April 29									30		30	17	22		1		11				5
Family TROCHILIDAE																						
Ruby-throated Hummingbird	May 15																10	9	13	16	17	13

Digitized by Google

Family TYRANNIDAE																						
Kingbird	May 3								29	30				21				10	6	10		5
Crested Flycatcher	May 3										30			21	29		3			11		
Phoebe	March 27		20			18	15	17	7		17	1										
Wood Pewee	May 9								27					28			19	10	11	16		12
Yellow-bellied Flycatcher	May 4											29									1	12
Least Flycatcher	May 4									30	20	30		21		12				6		9
Family CORVIDAE																						
Blue Jay	Resident																					
Crow	March 11		9	26	12	3	6	8														
Family ICTERIDAE																						
Bobolink	May 9																14		8	4	12	8
Cowbird	March 23		16			19	1	14	4		10	1										
Yellow-headed Blackbird	May 17																			17		
Red-winged Blackbird	April 5					22	26	17				18					7					
Meadowlark	March 17		14		27	19	10	13														
Orchard Oriole	May 8																15	6	11	10	2	5
Baltimore Oriole	May 3									30								2	6	4	2	5
Rusty Blackbird	March 31							17				19	5									
Bronzed Grackle	March 20	29	14	20	24	23	10	14														
Family FRINGILLIDAE																						
Purple Finch	April 16									24		15		11	15							
American Goldfinch	May 2								25		28		30				6		7		2	7
Vesper Sparrow	April 14											22		18	1							
Savanna Sparrow	March 19							19														
Henslow Sparrow	April 24									29	26	21	30	24	15							
White-crowned Sparrow	April 23.						11				29		20		29	5	9		2			
White-throated Sparrow	April 13						24	21	26	13	19	21	17									
Tree Sparrow	March 23						7	13				6	5									
Chipping Sparrow	April 12								15	13	18	8	8	13	12							
Field Sparrow	April 3						11	19			23	17	5									
Junco	March 16	28	9	14	23	18	8	13														
Song Sparrow	March 20		15	19	27	18	8	12	8													
Lincoln Sparrow	May 14																		10	23	12	10
Swamp Sparrow	April 14						11				20	21	18		12		7					

	Average First Appearance	MARCH							APRIL							MAY						
		1897	1898	1899	1900	1901	1902	1903	1897	1898	1899	1900	1901	1902	1903	1897	1898	1899	1900	1901	1902	1903
Family FRINGILLIDAE—Continued																						
Fox Sparrow	March 23		30	14		18	10	17	8			5										
Towhee	March 27		17			25	10	18	7		11	7										
Cardinal	April 11	28										25										
Rose-breasted Grosbeak	May 7													28					10	11	4	9
Indigo Bunting	May 11																14	10	21	10	2	6
Dickcissel	May 17																			16	18	
Family TANAGRIDAE																						
Scarlet Tanager	May 5																14	4	2	5	3	5
Summer Tanager	May 4																	4				
Family HIRUNDINIDAE																						
Purple Martin	May 6												27	23	17		25	30				
Barn Swallow	April 21								18		23	26	10	18	24		1					
Tree Swallow	April 11						11				12	18	12		1		11					
Family BOMBYCILLIDAE																						
Cedar Waxwing	April 16			27						11				24	5					16		
Family LANIIDAE																						
Loggerhead Shrike	March 13		14		9	3	9	1			11											
Family VIREONIDAE																						
Red-eyed Vireo	May 5										29						10		12	4	2	4
Warbling Vireo	May 8								27								15	12	6	11	7	5
Yellow-throated Vireo	May 3													28			1		12	5		12
Blue-headed Vireo	May 8																		17	5	2	8
Family MNIOTILTIDAE																						
Black and White Creeper	April 28.								28	30	21	26		28	27					4		
Prothonotary Warbler	May 11													27				16				19

Digitized by Google

Blue Golden-winged Warbler	May 10																14		12	12	2	
Nashville Warbler	May 7																10	4	9	19	2	1
Tennessee Warbler	May 11																11					
Northern Parula Warbler	May 8																11			12	3	9
Cape May Warbler	May 5																13	1	2	5	2	9
Yellow Warbler	May 2									30	29	30			29	11				6	2	
Black-throated Blue Warbler	May 7															11	11	4	8	5	2	5
Myrtle Warbler	April 14								14	16	14	19	14	17	8							
Magnolia Warbler	May 2								26		27			21			11		8	6		5
Cerulean Warbler	May 12																		12			
Chestnut-sided Warbler	May 12															21	10	30	3	7	2	9
Bay-breasted Warbler	May 9													22			20	4	11	15		9
Black-poll Warbler	May 13																19	10	13	16	12	6
Blackburnian Warbler	May 4											28		28	30	12	10	3		10		
Black-throated Green Warbler	April 29								18		23	27	26	28			17					2
Pine Warbler	April 25											26	30	21	22							
Palm Warbler	April 24									29	20	27	26	21	22							
Oven-bird	May 5															11	1	3	5	5	1	6
Grinnell Water-Thrush	April 25									30	28	23	17	21	29							
Louisiana Water-Thrush	April 30													27								5
Connecticut Warbler	May 20																					20
Mourning Warbler	May 11																20	13	6	16	2	11
Maryland Yellow-throat	May 1								26		28						1		5	5	2	1
Yellow-breasted Chat	May 6																				2	11
Hooded Warbler	May 12																					12
Wilson Black-cap Warbler	May 14															21	6	22	10	16	3	17
Canadian Warbler	May 13																19	16	12	16	6	12
American Redstart	May 5										28					11	1		11	6	2	5
Family TROGLODYTIDAE																						
Mockingbird	April 29												29									
Catbird	April 25						26		26		28	30	28				1					5
Brown Thrasher	April 22								26	24	27	16	24	22	12							
House Wren	April 22								14		30		21	23	12		4					
Winter Wren	April 13					22			8	25	30			19	8							
Long-billed Marsh Wren	April 8							29		14		12										
Family CERTHIIDAE																						
Brown Creeper	March 31						21	18	7	7	6	2	4									

Digitized by Google

	Average First Appearance	MARCH							APRIL							MAY						
		1897	1898	189	1900	1901	1902	1903	1897	1898	1899	1900	1901	190'	1903	1897	1898	1899	1900	1901	1902	1903
Family PARIDAE																						
White-breased Nuthatch	April 1		29			18	31		7		11	2			2							
Red-breasted Nuthatch	April 28													22	28				3			
Chickadee	April 1				19					1			13									
Family SYLVIIDAE																						
Golden-crowned Kinglet	March 31		14				28	19	7		11	6	5									
Ruby-crowned Kinglet	April 8							19	14	1	13	7	17	13								
Blue-gray Gnatcatcher	April 23									25	22	18	28	22	25							
Family TURDIDAE																						
Wood Thrush	May 1								18					29			3	1		4		12
Wilson Thrush	April 16									10	12	7	11	26	8	11						
Gray-cheeked Thrush	May 16																				21	11
Olive-backed Thrush	May 1												30	22	29		1	4	8			
Hermit Thrush	April 4		31					19			11	7	5	13								
Robin	March 15	28	9	14	24	14	10	7														
Bluebird	March 18.	28	8		31	17	7	7			4											

Digitized by Google

Notes.

Digitized by Google

Table of Distribution and Occurrence.

The following table has been compiled in the hope that wherever the bird student may be in the northern states, east of the Mississippi river, he may have at hand for comparison the results of some reliable observer in the same or a nearby area. No better method of gaining a clear idea of the movements of birds can be suggested than to study carefully their distribution and occurrence within related areas. In order to do this understandingly, the main topographic features, taken as a whole or state by state, should be considered. The bird population of any region will be found to be closely related to its mountains, river basins, prairies, forests and wet or dry areas.

Attention is called to the meeting and intermingling of the Carolinian and Alleghenian faunal areas in these northeastern states. The teacher of geography as well as the bird student will find great interest in following out the limits of life conditions in these areas.

The table contains in addition to the 200 species already described, 148 species of frequent but rarer occurrence throughout the area covered. It should be said that considerable difficulty lies in the way of reducing to a few arbitrary symbols the notes of several ornithologists whose data are not equally complete and who may not use the same descriptive terms, or who may attach a different sigificance to the same terms. The authors nevertheless have earnestly tried to interpret correctly each authority cited and trust that the useful features of the result will outweigh its shortcomings.

The *varying dates* of the pubished lists for the different localities given should be kept constantly in mind in making

Digitized by Google

comparisons with new lists, because the bird population of any locality is as truly changeable, although for different reasons, as its human population.

The data given have been compiled from the following published lists:—

1. **Maine.** Knight, O. W., Birds of Maine. 1908.
2. **New Hampshire.** Allen, Glover M., A List of the Birds of New Hampshire. 1903.
3. **Vermont.** Perkins, G. H., and Howe, C. D., A Preliminary List of the Birds found in Vermont. 1901.
4. **Massachusetts.** Brewster, Wm., The Birds of the Cambridge Region of Massachusetts. 1906.
5. **Massachusetts.** Clarke, H. L., The Birds of Amherst and Vicinity, 2nd edition. 1906.
6. **Rhode Island.** Howe, R. H. Jr., and Sturtevant E., The Birds of Rhode Island. 1899. Supplement, 1903.
7. **Long Island.** Braislin, W. C., A List of the Birds of Long Island, N. Y. 1907.
8. **New York City.** Chapman, F. M., The Birds of the Vicinity of New York City. 1906.
9. **Pennsylvania and New Jersey.** Stone, Witmer, The Birds of Eastern Pennsylvania and New Jersey. 1894.
10. **Ohio.** Jones, Lynds., The Birds of Ohio. 1903.
11. **Indiana.** Butler, A. W., The Birds of Indiana. 1897.
12. **Illinois.** Woodruff, F. M., The Birds of the Chicago Area. 1907.
13. **Missouri.** Widmann, Otto, A Preliminary Catalog of the Birds of Missouri. 1907.

Digitized by Google

KEY TO TABLE OF DISTRIBUTION AND OCCURRENCE.

Column I Number

- A.....abundant.
- a.....common or very common.
- F.....frequent, rather common, not uncommon.
- f.....infrequent, not common, rather rare, uncommon.
- —.....rare.
- =.....very rare or casual.
- x.....accidental, out of normal range.
- i.....irregular, from year to year common, rare or absent.
- (blank)..no record.
- d.....doubtful, records scanty or incomplete.

Column II Status

- R.....Permanent Resident.(May be partially migratory, increasing in number during the spring or fall or both; or within the limits of a single state withdrawing largely to the north in summer and to the south in winter in search of special breeding and feeding areas.)
- R.....Permanent Resident most abundant in winter.
- r.....Winter Resident.
- P.....Winter Resident with a few in summer.
- s.....Summer Resident.
- S.....Summer Resident with a few in winter.
- V.....Winter Visitant.
- W.....Winter Visitant with a few in summer.
- v.....Summer Visitant.
- T.....Regular Transient, a spring and fall migrant.
- t.....Regular Transient, most abundant in fall.
- ɜ.....Regular Transient most abundant in spring.
- I.....Regular Fall Transient, not recorded in Spring.
- I.....Regular Transient with some wintering locally.
- II.....Regular Transient with some summering, or breeding locally.
- III.....Regular Transient with a few winter and summer.
- IV.....Regular Transient frequent or common in winter, infrequent or rare in summer.
- X.....Regular Transient frequent or common in summer, rare in winter.
- d.....doubtful, may occur or probably occurs.
- ?.....no record within recent years.(usually indicates species formerly common).
- D.....an introduced species.

Column III Distribution

- g.....general throughout area cited.
- L.....not general, confined mostly or entirely to particular regions within area cited.
- l.....general, but restricted to particular breeding environments. (r. g. swamps dry pasture, second growth, etc.,) or, general, but for no apparent reason absent from many localities throughout area cited, hence local.
- c.....found on the coast.
- C.....found largely on the coast, a few inland.
- ɔ.....found largely inland, a few on the coast.
- n.....inland.
- m.....confined mostly to mountains (High altitudes are equivalent to northern latitudes).
- ?.....distribution doubtful.
- i.....irregular, or erratic, likely to occur at any point. (Usually applies to species going in flocks).

Figures in any column indicate number of recorded instances of rare species.

Symbols are occasionally placed without reference to the column-title in order to express the data given.

Digitized by Google

		Maine	New Hampshire	Vermont	Mass. (Amherst)
2	Holboell Grebe	f V C	— I ɔ	1	2
3	Horned Grebe	f IV C	F t n	f v	
6	Pied-billed Grebe	F II C	f II L	F s	F T
7	Loon	f X ɔ	a III C	f s	i T —
9	Black-throated Loon	? V c			
11	Red-throated Loon	F IV C	— t ɔ	— T	
13	Puffin	F P c	V c		
27	Black Guillemot	F P C	V c		
31	Brunnich Murre	f r C	a r C	— v	
32	Razor-billed Auk	f V c	f V c		
34	Dovekie	f V C	f V C	= V	x
36	Pomarine Jaeger	f II c			
37	Parasitic Jaeger	f II c	v c		
40	Kittiwake	a P C	a r c		
42	Glaucous Gull	— V c	— V		
43	Iceland Gull	f V c			
47	Great Black-backed Gull	F P c	a r c		
51a	Herring Gull	a X C	a I C	f s L	—
54	Ring-billed Gull	F I c	T c		
58	Laughing Gull	f s c		— s	
59	Franklin Gull				
60	Bonaparte Gull	a I C	f I C	— s	
64	Caspian Tern	— T c	= v		
69	Forster Tern				
70	Common Tern	a s c	II C		
72	Roseate Tern	— v c	?		
74	Least Tern	= ? 1	=		
75	Sooty Tern	= x 1	1 x		
77	Black Tern	= v c	= v C		
80	Black Skimmer	= ?			
88	Cory Shearwater				
89	Greater Shearwater	f c	1 c		
94	Sooty Shearwater	f v c			
106	Leach Petrel	F S C	a v C	= v	
109	Wilson Petrel	— v C	a v c		
117	Gannet	F I c	T c		
119	Cormorant	F r C		— T	
120	Double-crested Cormorant	F IV C	T C		
125	American White Pelican	= ?	?		
129	American Merganser	F III C	F III n	F S g	F T
130	Red-breasted Merganser	F III C	a I C	=	1
131	Hooded Merganser	f II C	f II L	— s	f T
132	Mallard	f I C	— T n	— T	
133	Black Duck	a II ɔ	a II n	a s	— T
135	Gadwall	d T c	?	1	
136	Widgeon			=	

Digitized by Google

Mass. (Cambridge)	Rhode Island	Long Island	New York City	Penn. and New Jersey	Ohio	Indiana	Illinois	Mo.
f I	i V C	r	f I	— V c	i I —	— T L	— V c	= ?
f t	ɹ r C	a I	i I ɔ	i I ɔ	f I	F III	F II	F T L
a t	f s n	f III n	a I	F III ɔ	F II g	a II 1	a s g	F II g
— T	ɹ IV C	a V	a I	a III C	f II c	F III	F R	f T g
		1 =	1		= V			=
= t	f III C	F T c	F I	— I c	— T	— I	— I	3 T =
		= V	1 L	= V				
	i V =	1	2 L	= V c				
— V t	i V C	F V L	F V L	f V C		= x		
	— V c	V c	F V L	f V c				
i V	i V C	f V c	F V L	i V c				
	f v c	f T c	f t c	= I C	3 T —	= V c	=	
	1	F T	f t c	= c	— T			= T
	f V c	a V	a T r	= V C			= V	1 x
— V	2	— V	f L			= W c	=	
— V		= V	1 L	= V		f V c		
a r	F r C	a r	F r C	— V c	= V			
A r	A P C	A IV	a IV C	A r C	a IV C	a I	a r C	F I 1
	f V	a III	f III C	a r C	d	F I 1	a r	a T
	1 =	— s	— s L	f II C				
						— T		— T L
— ɹ	a IV C	F V	F I	F I C	a T 1	a I	a T	F. T L
	3 v	— T	f t	= v C	= v		f t	= v
		2 x = L	— x	— v c	= v	A T C	f T	f T g
= T	a s L	a s L	a s L	f II C	a II 1	a II	A T	— T
	f v	f s L	— s	= v C	= v C	= x		
	— v C	— T	— T	— II C	— T g	— II 1	= v 1	— s
	x	= ?	2 x	= v C				
	i T c	a T	i T	t t C	F s L	F II 1	a s 1	f II 1
		— v c	= v c	— s c				
	f v c	v c	f T c					
	c	i v c	i v c	= v c				
f	c	= v c	f v c	— v c				
— t	f T C	— T	f T	— i C				
f	T C	a s c	F s c	F v c				
	f T c	F t c	T c	— T c				
	f r L	— T	—	— T c				
x t	a T C	a T v	a T	a III C	— T L	F 1 1	— t	a T L
		2	2	?	— T L	— T	= T	a II
f V	f V ɔ	f V	f r C	F III ɔ	f r C	a I	I c	a I
f t	a IV C	A IV	a T L	a III C	— I C	f I g	P C	— I
f t	F I	F III t	F I	f I ɔ	F II 1	A IV 1	a III	F III 1
f t	f V C	f T	i I	F III ɔ	A III 1	A III 1	a III	a IV
a II	A P	A III	III	a IV C	F II	f III	F T	— T
	?	=	= T	— T ɔ	F T	— T	— II	F T
		=	= x c	x T C	— T	— x	= ɹ	1 =

Digitized by Google

		Maine			New Hampshire			Vermont			Mass. (Amherst)		
137	Baldpate	f	T	C	—	T	c	—	T				
139	Green-winged Teal	f	t	C	—	T	n	—	s		—	T	
140	Blue-winged Teal	F	t	C	—	T	n	f	s	L	f	T	
142	Shoveller	—	T	c	=	T		=	T		=		
143	Pintail	f	T	c	f	T		—	T		1		
144	Wood Duck	—	II	l	f	II	1	F	R	T	F	II	
146	Redhead	f	T	C	—	t		f	T	L			
147	Canvas-back	—	T					=	T	L			
148.	American Scaup Duck	F	I	c	—	T	C	F	T	L			
149	Lesser Scaup Duck	f	I	c	—	T	n	2					
150	Ring-necked Duck	=	T	c	=	T							
151	American Golden-eye	a	IV	C	a	III	C	a		1			
152	Barrow Golden-eye	f	V	c	1		n	1					
153	Buffie-head	F	I	C	f	T	C	—	V				
154	Old-squaw	a	P	C	a	I	C	—		L			
155	Harlequin Duck	—	V	l	—	V	c						
160	American Eider	F	P	c	F	r	c						
162	King Eider	—	r	c	1								
163	American Scoter	F	IV	C	—	IV	C	1					
165	White-winged Scoter	a	IV	C	a	IV	C	1					
166	Surf Scoter	F	IV	C	—	IV	C	1					
167	Ruddy Duck	f	II	C	—	II			T				
169	Lesser Snow Goose	=	T		i	t	—	=		1			
169a	Greater Snow Goose	=	T	c									
169.1	Blue Goose				1	x							
171a	American White-fronted Goose							—		L			
172	Canada Goose	a	T		a	T	n	a	T		a	T	
172a	Hutchins Goose	=	T										
173	Brant	F	T	C	f	T	C	—	T			x	
180	Whistling Swan				=	x		=					
181	Trumpeter Swan												
184	White Ibis												
186	Glossy Ibis				1	v	x						
188	Wood Ibis	1	x						x				
190	American Bittern	F	s		F	s	n	F	s	l	f	s	
191	Least Bittern	—	s	1	—	s	L	—	s	l	=		
194	Great Blue Heron	F	s	ɔ	F	II	n	f	s	l	i	t	—
196	American Egret	=	v	5	1	v	x				1	x	
197	Snowy Heron										1	x	
200	Little Blue Heron	=	v	4	1	v	x						
201	Green Heron	f	s	1	f	s	L	F	s		a	s	
202	Black-crowned Night Heron	f	s	C	F	s	C	—	s	1	i	v	—
203	Yellow-crowned Night Heron	=	v	2									
204	Whooping Crane								?				
206	Sandhill Crane					?		1	x				
208	King Rail	4	x	C									
211	Clapper Rail	3	x		1	x	c	=	x			x	
212	Virginia Rail	f	II		—	s	L	F	s		f	s	
214	Sora	f	II	ɔ	f	II	L	—	s		i	s	
215	Yellow Rail	—	II	c	1			=			1	=	
216	Black Rail												

Digitized by Google

Mass. (Cambridge)	Rhode Island	Long Island	New York City	Penn. and New Jersey	Ohio	Indiana	Illinois	Mo.
— t	— V C	F IV t	i T	F t	F T 1	a II L	a II	a T
— T	f I ɔ	f t	f I	F II ɔ	F T	a I ʇ	a III	a T g
f t	F II 1	f T	a t	a T ɔ	F T 1	a II 1	a s	a II
= T	—	— T	i T —	T	— T	F III	a II	F IV
f T	f T c	F I	a T	—	a I	A ʇ	a T	a T g
f T	f II 1	— s	F II	F X n	f II	f III	F T	F II
f t	— I	F T L	F I L	f I ɔ	F . T	a T 1	a T	F ʇ
= T	i T =	— T	f T	— T L	— T	f T	f T	i T
	F I ɔ	A IV	A I	a I g	f II	— T	d	i I
F t	F T C	a I	a I	a V C	F II	A II	A II	a IV
= t	=	=	i T =	f I ɔ	— T	F T	A T	a T
a I	a I C	a V	f I 1	a I C	F I	a I	a r	F I
		1			3 V	— V	— V	— V
f t	F I C	F V	F I	a r C	a T	a I	f I ɔ	F I
f t	A r c	A V	a r	A r C	i r	a r C	a r	i V
	— V	= V c	= V c	= V c				— V
	a I C	— V c	— V	= V c	= V		= r	
= V	i V C	— V	V L	= V c	= V			1
— t	F r C	a W c	F I	a r C	4 V		= V	— T
f t	a P C	a V c	a I c	F V c	— I	— V C	— r	— V 1
f t	F P C	a W c	? I c	a r C		— r C	= V	— V
F t	F II C	i V 1	F T	F I ɔ	F T	f T 1	F T	F T g
	i V —			= V m	—	— ʇ	a T	F T g
	? ?	— T	i T	— V n	—	— T	?	d T
	1	=		= V n	f t	— T	a T	F ʇ
		=	4 T =	= V C	— I	— T	a T	F T
F T	a T	a I	a T	a I C	F I L	a III	a T	a III
		? —	— T		=	— I	— T	F I
	F I C	a T	a T	a I C		x	?	
x t	1 ?	— T	= T	— I C	— T	f I	— T	— ʇ
					= T	— T		— ʇ
		= ?	2 x	= x C		— v		2
1 ^		= x 2	2 x	= x n				1 v
	2 x	= x		= x n	1 x	i v L		f v L
a II	a II	a T	f s	F II 1	F II 1	F III 1	a s 1	a II g
F s 1	a s	a s	a s 1	F s C	f II 1	f II 1	a s 1	a s l
f I	a T C	— S	a II 1	a X ɔ	F s 1	a X 1	F s	F s g
	6 v =	— v	i v —	— v C	= v	— II	— v	= ? L
		= v	= v	= v C	= v	— II L	?	= ? L
v x	= v	— T	— v	— v ɔ	= s	F s L		F v L
a s	a s	a s	a s	a s g	a s	a s g	— s	a s g
f t 1	a S 1	a S	a s 1	c s ɔ	a s 1	F II L	A v	f s L
x	2 v =	= 3	1	= v ɔ		a s L		— s 1
				?	= T L	— T	= T	= ʇ
					— II	F II 1	= T	= T
=	3	— d	— s 1	f s n	F s C	a II 1	a s	F s 1
		a S L	a S c	A X C				
A s 1	F II	F s	a S 1	a II ɔ	a s C	F ʇ s	a s	F ʇ g
A s 1	a II	a II	a II	A II ɔ	F s g	a II	a s	a II g
— t	1	— s d	d t	— t n	f s	— II 1	F s	— ʇ
? d	4	— s d	— s	= t n	= d	— s 1	= s	— d

Digitized by Google

		Maine			New Hampshire			Vermont			Mass. (Amherst		
218	Purple Gallinule	3	x		2		C						
219	Florida Gallinule	—	T		1			=	s				
221	American Coot	f	t	C	—	t	L	—	T		i	T	—
222	Red Phalarope	—	II	c	a	T	C	=					
223	Northern Phalarope	F	I	C	A	I	C						
224	Wilson Phalarope	2	x		—	T	c						
225	American Avocet	1	x										
228	American Woodcock	a	II		f	II	1	F	T		f	s	
230	Wilson Snipe	a	II		f	t	C	f	s		i	t	—
231	Dowitcher	f	?	c	F	t	c						
232	Long-billed Dowitcher												
233	Stilt Sandpiper	—	T	c	—	t	c						
234	Knot	f	T	c	f	T	c						
235	Purple Sandpiper	F	V	c	2			—	v				
239	Pectoral Sandpiper	a	T	C	F	t	C	—	t				
240	White-rumped Sandpiper	F	t	C	—	T	C	1					
241	Baird Sandpiper	—	T		—	t	C						
242	Least Sandpiper	a	T	C	a	T	C	—	T		—	?	
243a	Red-backed Sandpiper	f	t	c	—	T	c						
246	Semipalmated Sandpiper	a	t	C	a	T	C	—	T				
247	Western Sandpiper				1								
248	Sanderling	F	t	C	a	T	c	—	v				
249	Marbled Godwit	4	x	c	1	x	c						
251	Hudsonian Godwit	—	T	c	1	t	c	—					
254	Greater Yellow-legs	F	T	C	a	T	C		T	1			
255	Yellow-legs	f	t		a	t	C	—			i	T	—
256	Solitary Sandpiper	a	II	1	a	II	3	F	II		i	T	—
258	Willet	=	x		i	t	c	1					
258a	Western Willet												
261	Bartramian Sandpiper	—	II	n	—	II	L	—	II		—	II	
262	Buff-breasted Sandpiper	—	T	1	—	t	c	—	T				
263	Spotted Sandpiper	a	s		F	s	g	a	s		a	s	
264	Long-billed Curlew	=	T	c	x	t	c						
265	Hudsonian Curlew	—	T	C	—	T	c					x	
266	Esquimaux Curlew	=	t										
270	Black-bellied Plover	F	t	C		T	c	2	T				
272	American Golden Plover	f	t	c	—	t	c	=	T				
273	Killdeer	—	T	c	=	?	C	—	II		i	T	—
274	Semipalmated Plover	a	T	C	a	T	C	—		1			
277	Piping Plover	—	T			T	c				1	T	
283	Turnstone	F	t	C	f	T	c	1					
289	Bobwhite	—	R	L	—	R	L	—	D		f	R	
300	Ruffed Grouse	a	R	*	F	R	*	a	R	g	a	R	
305	Prairie Hen					D	?						
310	Wild Turkey		?			?			?				
315	Passenger Pigeon		?			?			?			?	
316	Mourning Dove	—	v	L	F	s	L	=			a	s	

*Canadian Ruffed Grouse.

Digitized by Google

Mass. (Cambridge)	Rhode Island	Long Island	New York City	Penn. and New Jersey	Ohio	Indiana	Illinois	Mc.
	7 x	— x	2 x	= x	7 x	— ɿ	1 x	2 x
f s 1	f II 1	— s 1	— T 1	f t n	F s g	a II 1	a s	— s L
i T	F T ɔ	a II	F II 1	f II ɔ	f X	a III	a s	a II g
— ɿ	f T	— T c	i T —	= T c		= T	?	=
— ɿ	F T	f T c	F T	— T C	3 =	— T	— T	=
	? t 4	=	i T =	= T c	— T	— II L	a s 1	— T
		= v 2	= ? x	= v c	=	— T 1	?	— T
f II	F II	— S	F s T	F III g	F s 1	a S 1	— s	F X L
a T	a I ɔ	a I	F III	a III g	a I	A III	a T	a T g
	F T	a T	a T	a T C	= T	— t	= T	
	1	— T	— T c	— t C	= T	— T	= T	— T 1
	F t	i T	f t	= T C	= d	— t	f t	— ɿ
	F T	f T	a T	f T c	1	— T c	— t	x
x t	a r L	— V	— r	= V C	1		= T	1 ?
— t	A T C	a t	a t	F t C	a T g	a T	A T ɔ	a T
	f T	F T	F T	f T C	d T C		= T	— T
	7 t	= T	3 x c		d t	— t 2	— T	F T
a T	A T C	a T	a T c	A T C	F T	F t	A T	F T
— t	F T C	a T	a t c	A I C	f t	F T C	a T ɔ	— T
f t	A T C	A T	A t c	A T C	F T	f T	a T	a T
	f T	f T		= T C				=
— t	a T c	a T	A T c	A I C	F T C	F t c	F t c	— T
	= T C	= T	i T —	— T C	— T	— T		— T
= t	— T	— T	i T	— t C	— T	= T	— T	— T
a T	a T C	a T	a T	a T C	A T g	F T L	a T	F T g
— T	a t C	a t	A t	a T C	F T g	a t s	a T	a T
a T	f II C	F T	F T ɔ	F II ɔ	F II	a II L	a T	a T g
	— T c	f T c	— T	— II C	= ?	— T		— T
			d			— T	f t	— T
— t	f T 1	— s	f II	F II 1	F s	F II 1	a s n	f II 1
	f T	= T	= t	= t c	— T	— T	= T	— T
a s	A s g	a s	a s	A s g	a s	a s g	A s C	a II
	— T	= T	i T —	= T C	— T	— T	= T	— T
	f T c	F T	F T	a T C	= T	= T	= T	— T
	= T	= ?	1 ? =	= ? c	1 d	— T	?	— ɿ
	F T	a T	a t	a T C	— T	— T	— T	— T
	f T	F T	a t	i t ɔ	— T	F T	a T	a ɿ
— II	f II g	f III	F II	a X ɔ	a X	a S	a S	F II
f t	A T c	A T	a T c	a T C	F t	f T C	a T	F T 1
	F II c	— II	F II 1	f III C	— T	— T	— s	— T *
	a T c	F T c	a T	a T C	F ɿ C	— T C	— T c	— T
— R	a R †	a R	F R L	f R	F R g	F R	— R	f R
— R	a R L	F R	f R 1	— R L	f R 1	F R 1	= ?	— R 1
						— R L	— R	— R L
				= R m	?	R ? g	— 1	= R L
?		?	= ?	= ?	= T	— X	?	?
— T	F s L	F s	a S	a S	a X g	a S	a s	a S

*Belted Piping Plover.
†Reservations established and stocked.

Digitized by Google

		Maine			New Hampshire			Vermont			Mass. (Amherst)	
325	Turkey Vulture	8	x		=	x	C	2	x			
331	Marsh Hawk	F	II	1	f	s	1	F			a	s
332	Sharp-shinned Hawk	F	X	1	a	III	1	F	s		a	T
333	Cooper Hawk	f	II		F	II		a	s		a	s
334	American Goshawk	F	W		i	W		—	W		F	V
337	Red-tailed Hawk	f	III		f	R	L	a	R		a	R
339	Red-shouldered Hawk	a	II		—	R	L	a	R		a	R
343	Broad-winged Hawk	F	II	g	F	s	m	—	s		—	
347a	American Rough-legged Hawk	—	V		—	V	ɔ	—			a	r
349	Golden Eagle	=	s		=	R		=			=	x
352	Bald Eagle	—	s	C	f	S	1	f	R	1	—	x
354a	Gyrfalcon	4	V									
356	Duck Hawk	f	II	C	—	s	1	—	R	1	=	s
357	Pigeon Hawk	f	II		f	T	n	f	III		i	I
360	American Sparrow Hawk	F	II	g	f	II	L	a	R		a	s
364	American Osprey	F	s	C	F	II	C	—	R		a	1
365	American Barn Owl							—		L		
366	American Long-eared Owl	f	R	g	—	R	L	F	R		f	R
367	Short-eared Owl	—	R	l	—	T	n	f	R		—	V
368	Barred Owl	F	R		F	R		F	R		f	R
370	Great Gray Owl	i	V		—	V	n	=	V		=	V
371	Richardson Owl	—	V		—	V	n	—	V			
372	Saw-whet Owl	f	R	1	f	R	g	F		1	=	
373	Screech Owl	f	R	g	f	R	l	a	R		a	R
375	Great-horned Owl	f	R	g	f	R	l	F	R		a	R
376	Snowy Owl	i	V		i	V	ɔ	i	V		i	V
377a	American Hawk Owl	i	V		i	V	—	—	R	L	=	V
387	Yellow-billed Cuckoo	=	s	L	=	s	L	—	s		f	s
388	Black-billed Cuckoo	a	s	g	a	s	l	f	s		a	s
390	Belted Kingfisher	a	s		F	S	n	a	s		f	s
393	Hairy Woodpecker	F	R	g	F	R	m	F	R		f	R
394c	Downy Woodpecker	a	R	g	a	R		a	R		a	R
400	Arctic Three-toed Woodpecker	f	R	L	—	R	m	—	W	1	=	V
402	Yellow-bellied Sapsucker	F	II	g	F	II		a	s		—	T
405	Pileated Woodpecker											?
405a	Northern Pileated Woodpecker	f	R	L	—	R	m	—				?
406	Red-headed Woodpecker	—	II	L	—	v		—	s		i	v
409	Red-bellied Woodpecker											
412a	Northern Flicker	a	X	g	F	S	ɔ	a			a	s
417	Whip-poor-will	F	s		a	s	1	f	s	i	F	s
420	Nighthawk	a	s		a	II	1	a	s		a	s
423	Chimney Swift	a	s		a	s	1	a	s		A	s
428	Ruby-throated Humming-bird	a	s	g	a	II		a	s		a	s
444	Kingbird	a	s	g	F	s		a	s		A	s
452	Crested Flycatcher	f	s	L	—	s	1	a	s		f	s
456	Phoebe	a	s		F	s		F	s		A	s
459	Olive-sided Flycatcher	F	s	1	—	s		a	s			

Digitized by Google

Mass. (Cambridge)	Rhode Island	Long Island	New York City	Penn. and New Jersey	Ohio	Indiana	Illinois	Mo.
v	= v 4	— v	f v L	a S 1	f S	f R 1	— v.	F X g
T	F II	a III	a X	F R C	F r	F III 1	a III	F II L
T	f X	a III	f S	a R g	F R	F III	i II	F II g
X	a X	F III	F S	a R	F R	F X	F s	F II g
V	i V —	— V	F V	a V	f V	— V	i T	i T
I	F W	a R	a R	a R 1	f R g	a R	a R	f III
I	a R	a R	a R	F R	F R	a R	a R	a III g
T	f II	— R	f s	— R	f X	f III	F II	F II L
T	F I	a V	— r 5 L	f V L	f V	f r	i r	i r
	= 5	=	= r 5	i V	— V	— V	= V	— V
i	f i 1	— s 1	f R L	f III 1	f R 1	f R 1	— R	= s
	= 2		1 ?					
V T	— T C	a t	F II C	f III C	f ? g	f R 1	= T	— II L
I	a T r	a t r	a T C	— V C	— IV	f I	f III	f T
X	f s	a X	A III	A V g	a R	a X	a s	a X
T	a s L	A s L	A s L	a II C	f III 1	F III	— T C	= II 1
	= 5	— R	— R	f R L	f R	f R 1	=	— R L
R	f R	a II	f R	f R	f R g	F R	f R 1	F IV
T	f I	a II	a T	F R 1	f I g	f R 1	f s	F IV 1
R	f R	— T	a R	f W 1	F R g	F R	f R 1	F R g
V	= V		1 V	= V	2	V x	= V 2	
V	i V =					= V 3		
r	f V	— R	F r L	— III m	f R	F R 1	f IV 1	f W
R	a R	f R	a R	A R g	a R	f R 1	f R 1	a R g
V	— II	— R	f R	f R f	f R V	a R	= R	f R g
V —	i V f	f V	i V	i V C	— V C	a V f	i V	— V
V	1	1	1	= V	= V	x	2 1 ?	
s	a s	a s	a s	a s g	a s g	a s L	a s	a s g
s	F s	a s	a s	F s L	F s g	a s L	F s	f s L
X	a S	a s	a s	a X g	a X 1	a S	a s	F s g
V	f W — R L	— R	f R	a R g	F R	F R	a R 1	F R g
R	a IV	f R	a R	A R g	a R	a R	a III	a R g
		1			= V		= V	
I	F T	F T	a T	a III	a III	a IV	a T	a IV
		= ?		= R L		— R 1		
			= ?		— R L		= W 1	— R L
III f	i II	— s	F III 1	f s i	A S	a X	a S	A III g
	1 x	= v	— i	— V	F R	F R	— T	F R g
X	A R	A S	A X t	A S *	A III	a IV	a R	A X g
II	a s L	a II L	a s L	F II L	a s 1	a s	— s	a II
II	f II l	a s L	F s L	a s l	a s 1	A t s	a II	a II 1
s	a s	a s	A s	A s g	A s	A s	a s	A s g
II	a s	f s	a s	a s l	a s	a s	a s	a s
s	A s	a s	a s	a s g	a s	a s	a s	a II
s	f s	a s	a s	a s g	F s	a s	F II	a s g
II	a s 1	a s	a s	A s	a s	a s	a s	F s g
T	— T	f t	F t	= T	— T L	f T	= T	F T

*Southern Flicker

Digitized by Google

		Maine			New Hampshire			Vermont			Mass. (Amherst)		
461	Wood Pewee	F	s		F	s		a	s		a	s	
463	Yellow-bellied Flycatcher	F	II	L	a	s	m	—	s	L	i	T	f
465	Green-crested Flycatcher												
466a	(Alder Flycatcher)	a	s	l	f	s	l						
466	Traill Flycatcher							—	s	l	f	II	
467	Least Flycatcher	a	s	g	a	s	l	a	s		a	s	
473	Skylark												
474	Horned Lark	F	r	C	a	I	C	F	II				
474b	Prairie Horned Lark	f	II	n	—	II	L	1					
477	Blue Jay	a	R			R	g	a	R		A	R	
484	Canada Jay	F	R	L	F	R	m	f	W	L			
486	American Raven							=		l			
486a	Northern Raven	f	R	C		x					1	x	
488	American Crow	a	X		a	X		a	R		A	R	
490	Fish Crow							1					
493	Starling	1	x										
494	Bobolink	a	s	l	a	II	l	A	s		a	s	
495	Cowbird	a	s		f	S	L	a	s		a	s	
497	Yellow-headed Blackbird	1	x										
498	Red-winged Blackbird	a	s	l	F	s	l	A	s		a	s	
501	Meadowlark	f	S	L	f	S	l	f	s	l	a	s	
506	Orchard Oriole	3	x					—	s		i	s	—
507	Baltimore Oriole	a	s	L	f	s	l	a	s		a	s	
509	Rusty Blackbird	F	II		a	II	L	f	T		—	T	
511	Purple Grackle							F	s	d	?	s	
511b	Bronzed Grackle	a	s		f	s	l	f	s	d	a	s	
514	Evening Grosbeak	i	V	—	x	V					=	x	
515	Pine Grosbeak	i	W	g	f	R	*	i	V		i	V	F
517	Purple Finch	a	S	g	F	S		A	s		a	s	
521	American Crossbill	i	R	L	i	R			W		i	I	
522	White-winged Crossbill	i	R	L	f	R	m		W		—	V	
527a	Hoary Redpoll	1	V										
528	Redpoll	i	P	g	i	V		i	V		i	V	
528a	Greater Redpoll	i	V	C									
529	American Goldfinch	a	S	g	a	R		a	s		a	R	
533	Pine Siskin	i	R	i	a	I	m	—	V		i	V	—
534	Snowflake	a	V	C	F	I	C	i	V		i	V	—
536	Lapland Longspur	—	V	C				—	V				
537	Smith Longspur												
540	Vesper Sparrow	a	s		F	s		a	s		a	s	
501	Ipswich Sparrow	f	T	c									
542a	Savanna Sparrow	a	s	C	a	s		f	s		f	T	
546	Grasshopper Sparrow	—	s	L	—	s	L	f	s				
547	Henslow Sparrow				f	s	l	—	s		f	s	
548	Leconte Sparrow												
549	Sharp-tailed Sparrow	—	s	c		s	c	1					
549.1	Nelson Sparrow	—	T	d									
549a	Acadian Sharp-tailed Sparrow	f	T	c									
550	Seaside Sparrow	1	v										
552	Lark Sparrow												
553	Harris Sparrow												

*Canadian Pine Grosbeak.

Digitized by Google

Mass. (Cambridge)	Rhode Island	Long Island	New York City	Penn. and New Jersey	Ohio	Indiana	Illinois	Mo.
a II F T i II — a s	a s f T a s	a s f T f s 1 1 f s 1	a s F a t s f II L a s	A s — II L a s 1 — T F II 1	a s ? T F s 1 F II a II 1	a s — II a II d a s 1 F II L	a s F F T 1 F F s F II f s	a F II g F a T L F 1 s 1 F F II L
F I = T	a I = 1	— R L A V — d	f D L a r C — S	A r C — S L	a r L a R	? V a IV	— r a R	d V F R g
a R 1 2 = A III	a R A R	a R 2 = a R a X	a R t = A R a X L	a X g = V — R 1 A R g F III C	a R L = R L a R	a R — R A X	A R = V A R	a III g ? a R L
F s 1 a S A S F X 1 i s 1 A s a T — s L A S L	A s A S A s A R — s a T a s L	a R L a S 1 a s a S f s a T a s L a = T	D L a s L a S a II 1 a S a s F s a I 1 F s T	A II ɔ a S g A s n a R g a s g F S g F I a S	a II L a s a s 1 a X a s g F s a I a S	a s 1 A S f s L A S 1 F S f s a 1 a II	a s A a s 1 A a s a a S f a s A T a s	F II a X L a II L A X a X L a s g a I g a III g
i V = i V f a R F i — V i V a a V f i V R a F V V F 4 a s T — A II 1 f s = s ? s T 1 ? T ?	i V — a S a i I = i V f i V 1 A S a V ? T a s c a A II C A F s 1 3 ? c a s T ? T F s L	i V f F III a W — V = i V f a R F V f V a S a A V L A III 1 a T a S T L a s L f a	= V i a V a III i W F — V f i 2 V a F R A — r C a s a A r c a III L l a s l A T L A s T L f f s L A 2 x	i V x i V L — = 1 ? V = g F I 1 i F — V f V a R F V l L i V F V A II F II 1 l F s l d s L 1 d F s L	i V f i V — i W f i V — i V a V a X i V i V i T C a S g F III 1 a s a s f II L — T L F s	i V = i V i V i V i ? r a V a i i R i V A I a T a s F — — II 1 — s =	= V = a V a I i = I i V F X i V F T L F II l a II F a s L a II l f s g a T L

Digitized by Google

		Maine			New Hampshire			Vermont			Mass. (Amherst)		
554	White-crowned Sparrow	F	T		F	t		a	T		i	↕	
558	White-throated Sparrow	a	s		a	II		a	T		a	T	
559	Tree Sparrow	a	I		a	I			r		a	r	
560	Chipping Sparrow	a	s		a	s		A	s		A	s	
563	Field Sparrow	F	s	1	a	s	1	F	s		a	s	
567	Junco	A	X		A	X	1	a	III		a	r	
581	Song Sparrow	a	S	g	a	s		A	s		A	s	
583	Lincoln Sparrow	f	T		f	t		f	s		—	T	
584	Swamp Sparrow	a	II	1	F	II	L	—	s		F	s	1
585	Fox Sparrow	F	T		F	T			T		a	T	
587	Towhee	f	s	L	f	s	L	F	s	L	a	s	
593	Cardinal	1	x						x		=	x	
595	Rose-breasted Grosbeak	F	s	1	f	s	1	a	s		f	s	
598	Indigo Bunting	f	s	1	F	s		F	s		a	s	
604	Dickcissel	3	x										
608	Scarlet Tanager	f	s	1	F	s	1	f	s		F	s	
610	Summer Tanager	1	x					—	x				
611	Purple Martin	a	s	1	f	s	1	—	s	1	—	s	1
612	Cliff Swallow	a	s	1	a	II	1	a	s		a	s	
613	Barn Swallow	a	s		a	II		a	s		A	s	
614	Tree Swallow	F	s	1	a	II			s		a	s	
616	Bank Swallow	a	s	1	a	II	1	a	s		a	s	
617	Rough-winged Swallow							1			=		
618	Bohemian Waxwing	=	V		=	V		—	i				
619	Cedar Waxwing	a	S	i	a	X	i	F			a	s	
621	Northern Shrike	F	r	g	F	r	g	f	W		f	V	
622	Loggerhead Shrike				—	s	1	—			=	x	
622a	White-rumped Shrike							—	T				
622e	Migrant Shrike	f	s										
624	Red-eyed Vireo	a	s	g	a	s		a	s		A	s	
626	Philadelphia Vireo	—	II	1	—	II	L	f	s				
627	Warbling Vireo	F	s	L	f	s	1	F	s		F	s	
628	Yellow-throated Vireo	—	s	L	f	s	L	f	s	L	F	T	
629	Blue-headed Vireo	F	II	1	f	s		—	s		a	II	
631	White-eyed Vireo							—	s		=	s	1
636	Black and White Creeper	a	s	L	a	II		F	s		a	II	
637	Prothonotary Warbler	1	x					—	v	L	=	x	
639	Worm-eating Warbler							—	s		=	x	
641	Blue-winged Warbler							=	v	2			
642	Golden-winged Warbler				=	T		—	s		=		
645	Nashville Warbler	a	II	1	a	II		—	s		a	T	
646	Orange-crowned Warbler				1	x							
647	Tennessee Warbler	F	II		—	II	1	—	s				
648a	Northern Parula Warbler	F	s	1	a	II	1	a	s		a	T	
650	Cape May Warbler	—	II	1	—	II	1	—	s	2	i	↕	—
652	Yellow Warbler	a	s	1	F	s	1	a	s		a	s	
654	Black-throated Blue Warbler	F	II		F	s	1	F	s	L	a	T	
655	Myrtle Warbler	a	X		a	II		a	III		A	T	
657	Magnolia Warbler	a	II	1	F	II		f	s		f	T	
658	Cerulean Warbler										=	x	
659	Ches nut-sided Warbler	a	s	1	F	s	1	F	s		a	II	
660	Bay-breasted Warbler	—	II	1	f	II		f	T		i	T	—
661	Black-poll Warbler	a	II		a	II		F	II	m	f	T	

Mass. (Cambridge)	Rhode Island	Long Island	New York City	Penn. and New Jersey	Ohio	Indiana	Illinois	Mo.
f T	f T	— T	f T	— T	F T	a I	a T	a I g
a I	a I	A I	A I	a IV	a T	A T	A T	A I g
A r	A r	A V	A r	A r g	A r	a r	a r	a r
A s	A s	a s	A s	A S g	A s	a II	a s	a II g
a S	a S	a s	A S	A s I	a s 1	A S	a s	a X g
A I	a r	a V	A r	A W g	a r	a I	A I	A I g
A X	A S	A R	A S	A R	a X	a X	a s	a IV g
F T	f T	— T	— T	i T —	F T	— T	F s	F T g
A X 1	a II 1	a III	A S 1	a III 1	F II 1	a X	a s	a III g
a I	a T	a T	a I	a I	a T	a I	a T	a I g
a s 1	a s	a S	A S	a s 1	a X	a S	a s	a X I
f i		— s	F R L	F R 1	a R	a R	i s —	a R g
a s	a s L	f T	a s L	f II L	F II i	i II L	a s	a II L
a s	a s L	a s	a s	a s g	a s g	a s g	a s	A II g
? —	1	3 x	— i	= ?	a s L	A s 1	a s	a s I
F s	a s L	a s	a s	F II 1	a s	a s g	a s	F II 1
1 v	2 v =	= v	— i	= v	F s L	a s L	= v	a s L
— s 1	a II	a s 1	F s L	a II 1	F s 1	a s 1	a s	a s g
— s	f s	— s	a II t	f II C	a s 1	F s 1	a s	a II l
	A s	a s	a II t	A s g	A s g	A s	a s	a II g
F II	A II	A s 1	a II t	A II C	F s	i II 1	a II	ı II l
F II 1	a s 1	a s 1	a s 1	F II L	a s 1	A s 1	a s 1	a II 1
		— s	a s L	a s L	F s	f s 1	— ʇ	F s g
		?	= ? V	= V	?	i V	i V	= V
a III	a S	a R	a S	a R i	a R i	a III i	a S	F II i
a r	F V	f V	f V	— V	— r g	F V L	a r	F r
				— s 1	d	F X	a s	
					d			
= I	— T V	— T	— T		F S 1			f X 1
A s	A s	a s	A s	A s g	a s	A II	a s	a s g
— T	= T	— T	= T	= T	— T	— II L	— T	f T L
a s 1	a s	— s 1	a s 1	f s i	a s g	a II	f II	F s g
a s	F s	F s 1	a s	F II 1	F s l	a II	F II	a s g
F II	a II	f T	F T	F II L	F T 1	F T	f T	F T
— s 1	f s 1	a s	a s	a s 1	F s L	a s 1	— s	a s g
A II	a s	a s	a II	A II	a II	a II	a T	a II
	3 x =	2 x	3 x	= v	f s L	F s L	— v	a s
1 t =		— T	— s L	F s L	F s L	F s L		F s L
	= s 1	a s L	a s L	F s L	F II	F s 1	— T	F s l
F s	= s	— T	F t	— II	F II L	f II	f T	F II L
A II	F II	— T	F II L	F II	a T	F T	f T	a T L
— t	= T	x	= t 6	= T	— T	— i T	— T	a T L
i ʇ —	— T	f τ	f t	— t	a T	a t	a T	A T g
a T	a II L	A II 1	a II 1	A II 1	F ʇ	? II ?	a T *	F II *
— T	— T	— T	— ʇ	— T i	f t	i t	F T	— T
A s	a s	a s	a s	a s i	a s	A II	a s	a II g
F T	F T	a T	a T	a II	F T	F T	a T	— T l
A I	A I	a I	A I	A IV	i T	A I	A T	A I g
F T	F T	a T	a T	a II	F T	F T	A T	a T L
	= T 2	1 x	3 x	= T	a II 1	a II 1	— s 1	a s g
A s	a s	a T	a II 1	A II	a T 1	F II L	A T	F II l
f T	f T	— T	— T	f T	f t	F t	a T	f T L
A T	A t	a T	A T	A T	F T	i t	a T	a T

* Western Parula.

Digitized by Google

		Maine	New Hampshire	Vermont	Mass. (Amherst)
662	Blackburnian Warbler	F II 1	F s m	a s	f T
663	Yellow-throated Warbler				
663a	Sycamore Warbler				
667	Black-throated Green Warbler	a II	a II	a s	a T
670	Kirtland Warbler				
671	Pine Warbler	F II 1	F II 1	—	a T
672	Palm Warbler		f t	— T 2	
672a	Yellow Palm Warbler	a II 1	a ↕	— T	a T
673	Prairie Warbler		= s L		
674	Ovenbird	a s	a s	f s	a T
675	Water-Thrush	F s L	F II 1	— s 1	i ↕ —
675a	Grinnell Water-Thrush				
676	Louisiana Water-Thrush	1 x			= v
677	Kentucky Warbler				
678	Connecticut Warbler	— t L	— t	— s	= T
679	Mourning Warbler	f ↕ L	f s L	— ↕	= T
681	Maryland Yellow-Throat	a s	a s	a s	a s
683	Yellow-breasted Chat	3 x		= x	= v
684	Hooded Warbler	1 x		= s 1	
685	Wilson Warbler	f II	— II 1	— s	i ↕ —
686	Canadian Warbler	F II 1	a II m	— s	f T
687	American Redstart	a II	a II	F s	a II
697	American Pipit	F t	f t C	— T	= T
703	Mockingbird	d x		= s 1	
704	Catbird	a s L	F s 1	A s	A s
705	Brown Thrasher	F s L	F s L	f s 1	a s
718	Carolina Wren	— x	1 x c	= v L	
719	Bewick Wren		1 x		
721	House Wren	f s 1	f s 1	a s	f s
721b	Western House Wren				
722	Winter Wren	a S L	F X m	— II	i t —
724	Short-billed Marsh Wren	— s 1	— s L		
725	Long-billed Marsh Wren			= 1	
726	Brown Creeper	a R L	F R	f s	a IV
727	White-breasted Nuthatch	a R	a R	a R	A IV
728	Red-breasted Nuthatch	a R L	a R m	f IV	i V —
731	Tufted Titmouse	x		— R *	
735	Chickadee	a R g	a R	a R	A IV
736	Carolina Chickadee				
740	Hudsonian Chickadee	f	F R m		
748	Golden-crowned Kinglet	a R g	a III	a I	F V
749	Ruby-crowned Kinglet	F II	a T	a II	a T
751	Blue-gray Gnatcatcher	x			
755	Wood Thrush	— s L	— s	f s L	F s
756	Wilson Thrush	a s L	a s	F s	a s
756a	Willow Thrush				
757	Gray-cheeked Thrush	f T	— T	— s 1	
757a	Bicknell Thrush		a s m	— II	
758a	Olive-backed Thrush	a s L	F II 1	f s L	i T —
759b	Hermit Thrush	a s 1	F s	a s	a T
761	Robin	a S 1	A X	A S	A S
766	Bluebird	a II 1	a s	A s 1	a s

*No records cited. Entirely accidental in Vermont.

Digitized by Google

Mass. (Cambridge)			Rhode Island			Long Island			New York City			Penn. and New Jersey			Ohio			Indiana			Illinois			Mo.		
f	ɪ		f	T		F	T		F	t		a	II	1	F	T		a	T		a	T		—	T	g
						1	x		1			=	·													
															f	s	L	a	s	L		?		f	s	L
A	s		a	s		a	II			II	L	A	II		a	s	L	a	s	L	a	II		a	T	g
															F	T	1	a	T		2	=		1	ɪ	
a	S	1	a	s	1	a	s	L	F	II	L	a	s	1	d	II	d	=	II	1	—	T		F	II	L
f	t		—	ɪ			T		f	T					f	ɪ		F	ɪ		A	T		a	T	
a	T		a	s	1	F	T		a	T		A	I													
a	s	1	F	ɪ		F	s		F	II	L	A	s	1	—	II	d	—	II		=	v	ɪ	a	s	L
a	s		A	s		a	T		A	T		a	s	g	a	s	g	a	s		a	II		a	II	l
A	T		F	T		a	T		2	x		a	II		F	T	l	—	II		1	T	?	?	T	?
							x		a	s	L	=	T					?	T		A	T		a	T	g
			—	s		—	d	g	F	t	L	F	s	1	F	II		a	T	L	—	v		a	s	
a	t	1	=	d		=	t		f	T		f	t	L	f	T	L	a	s	L	=	II	ɪ	f	T	L
—	ɪ		f	—		F	T		—	s		=	ɪ		—	s		—	T		F	ɪ		—	s	
A	S		—	A		—	s		A	s		A	s	g	f	T	g	f	s		a	T		A	s	g
i	s	—	a	s	L	a	s	l	a	s		a	s	L	a	s	L	a	s	1	F	s		a	s	l
=			1		=	—	T		a	s	g	a	s	L	a	T		F	T		a	T		a	T	
a	ɪ		f	T		f	T		F	T		a	T		F	T										
a	II	ɪ	f	II		a	T		a	T		f	T		f	t										
A	s		a	s		a	s		A	s		A	II		a	s										
a	t		a	T		f	T		A	T	C	a	r	C	F	ɪ		A	I		F	T		F	T	
—	III		=	s		—	v		—	S		=	s		—	d		f	S		=			F		L
A	S		A	S		a	S		A	s		A	S	g	a	s	g	a	X		A	s		A	s	g
a	s		a	s		a	s		a	s		A	S	g	A	X		a	X		a	s		a	X	
—	v		=	s		—	R	1	F	III	L	a	R	l	a	R	L	A	S	L	A	R	1	a	R	L
												=	s	1	a	R	L	1	?	1	=	s	i	a	X	L
f	s	1	a	s	1	—	s		a	s		=	s	1	F	s		a	I	L	F	s		a		?
																		f	s	L	a	T		a	II	L
f	I		f	ɪ	r	f	I	t	F	r		f	IV	1	f	I		a	I	L	F	s		F	I	L
—	II	1	—	s		1			a	s	1	—	III	1	—	s		f	II	1	F	s	1	F	II	l
a	S	1	A	s	L	f	s	L	A	s	1	A	S	1	A	s	1	A	II	1	A	s	1			
a	I		a	I		a	I		a	I		A	IV		F	I		a	III		a	I	1	a	I	g
a	IV		F	IV		a	X		a	R		a	R		a	R	g	a	R		a	III		a	R	g
i	IV	f	F	I		f	V		F	ɪ	r	i	I		F	I		i	III	i	a	T		i	T	
						=			F	R	L	a	R	g	a	R	g	a	R	L	a	I		a	R	g
a	R		A	R		a	R		a	R	t	a	w	L	a	R	L	a	R	L	a	IV		a	R	L
									f	R	L	A	R	L	a	R	d	a	R	L	=			a	R	L
i	V	—	=	V																	=	V				
a	I		a	I		a	I		F	r		A	IV		a	I		A	I		a	T		a	I	
F	T		a	T		a	T		A	t		A	t		a	T		a	I		A	T		a	I	g
			=			=			f		L	—	s	L	F	s	l	a	s	1	F	II		a	s	l
a	s	1	a	s	1	a	s		A	s		F	s	g	a	s	g	a	s		a	s		a	?	g
A	s	1	A	s		f	s		a	s		a	II		F	II		—	II	1	a	T	1	f	T	g
			2	x														—	T	c	f	T		a	=	
f	T		F	T		a	T		a	T		F	T		f	T		f	T		A	T		a	T	
F	T		F	T		f	T		f	T		—	II											F	T	
a	T		F	T		a	T		a	T		a	X		a	T		a	T		A	T		A	T	g
a	I		a	T		a	T		A	I		A	III		a	T		a	T		A	s		a	I	g
A	X		A	S		A	S		A	X		a	X		A	X		A	X		a	s		F	I	g
A	X		a	S		f	s		A	X		a	X		a	X		a	X		a	s		a	X	g

Digitized by Google

Memo.

Digitized by Google

Index of Common Names

Bittern, American..... 19
Least..... 19
Blackbird, Red-winged..... 34
Rusty..... 35
Yellow-headed..... 34
Bluebird..... 52
Bobolink..... 33
Bob-white..... 52
Buffle-head..... 18
Bunting, Indigo..... 39
Cardinal..... 39
Catbird..... 49
Chat, Yellow-breasted..... 48
Chickadee..... 50
Coot, American..... 21
Cowbird..... 34
Creeper, Black and White..... 42
Brown..... 50
Crossbill, American..... 35
Crow..... 33
Fish..... 33
Cuckoo, Black-billed..... 29
Yellow-billed..... 29
Dickcissel..... 39
Dove, Mourning..... 25
Duck, American Scaup..... 17
Black..... 16
Lesser Scaup..... 17
Ruddy..... 19
Wood..... 17
Eagle, Bald..... 27
Finch, Purple..... 35
Flicker..... 30
Flycatcher, Crested..... 31
Green-crested..... 32
Least..... 32
Olive-sided..... 31
Traill..... 32
Yellow-bellied..... 32
Gallinule, Florida..... 21
Gnat-catcher, Blue-gray..... 51
Golden-eye, American..... 17
Goldfinch, American..... 36
Goose, Canada..... 19
Grackle, Bronzed..... 35
Purple..... 35
Grebe, Horned..... 13
Pied-billed..... 13
Grouse, Ruffed..... 25
Grosbeak, Rose-breasted..... 39
Gull, Herring..... 14
Bonaparte..... 14
Ring-billed..... 14
Hawk, American Sparrow..... 27
Broad-winged..... 27
Cooper..... 26
Marsh..... 26
Red-tailed..... 26
Red-shouldered..... 26
Sharp-shinned..... 26
Heron, Black-crowned Night..... 20
Great Blue..... 19
Green..... 20
Hummingbird, Ruby-throated..... 31
Jay, Blue..... 33
Junco..... 38
Killdeer..... 24
Kingbird..... 31
Kingfisher, Belted..... 29
Kinglet, Golden-crowned..... 51
Ruby-crowned..... 51
Lark, Horned..... 33
Prairie Horned..... 33
Loon..... 13
Mallard..... 15
Martin, Purple..... 40
Meadowlark..... 34
Merganser, American..... 15
Hooded..... 15
Red-breasted..... 15
Mockingbird..... 49
Nighthawk..... 30
Nuthatch, Red-breasted..... 50
White-breasted..... 50
Old-Squaw..... 18
Oriole, Baltimore..... 34
Orchard..... 34
Osprey, American..... 27
Oven-bird..... 46

Digitized by Google

Index of Common Names—Continued

Owl, American Long-eared..... 27
Barred..... 28
Great Horned..... 28
Saw-whet..... 28
Screech..... 28
Short-eared..... 28
Pewee, Wood..... 32
Phoebe..... 31
Pintail..... 16
Plover, American Golden..... 23
Semi-palmated..... 24
Pipit, American..... 48
Rail, Virginia..... 20
Red-head..... 17
Redpoll..... 35
Redstart, American..... 48
Robin..... 52
Sanderling..... 22
Sandpiper, Bartramian..... 23
Least..... 22
Pectoral..... 22
Semipalmated..... 22
Solitary..... 23
Spotted..... 23
Sapsucker, Yellow-bellied..... 30
Scoter, White-winged..... 18
Shrike, Loggerhead..... 41
Northern..... 41
Siskin, Pine..... 36
Sora..... 20
Snipe, Wilson..... 21
Snowflake..... 36
Sparrow, Chipping..... 38
Field..... 38
Fox..... 39
Grasshopper..... 37
Henslow..... 37
Lincoln..... 38
Savanna..... 36
Song..... 38
Swamp..... 38
Tree..... 37
Vesper..... 36
White-crowned..... 37
White-throated..... 37
Swallow, Bank..... 41
Barn..... 40
Cliff..... 40
Tree..... 40
Swift, Chimney..... 31
Tanager, Scarlet..... 40
Summer..... 40
Teal, Blue-winged..... 16
Green-winged..... 16
Tern, Common..... 14
Thrasher, Brown..... 49
Thrush, Gray-cheeked..... 51
Hermit..... 52
Olive-backed..... 52
Wilson..... 51
Wood..... 51
Titmouse, Tufted..... 50
Towhee..... 39
Vireo, Blue-headed..... 42
Red-eyed..... 41
Warbling..... 42
White-eyed..... 42
Yellow-throated..... 42
Warbler, Bay-breasted..... 45
Blackburnian..... 45
Black-poll..... 45
Black-throated, Blue..... 44
Black-throated, Green..... 45
Blue Golden-winged..... 43
Blue-winged..... 43
Canadian..... 48
Cape May..... 43
Cerulean..... 44
Chestnut-sided..... 44
Connecticut..... 47
Hooded..... 48
Magnolia..... 44
Mourning..... 47
Myrtle..... 44
Nashville..... 43
Palm..... 46
Northern Parula..... 43
Prairie..... 46
Pine..... 45
Prothonotary..... 42
Tennessee..... 43
Wilson, Black-cap..... 48
Yellow..... 44
Yellow Palm..... 46

Digitized by Google

Index of Common Names—Continued

Water-Thrush..... 46
Grinnell..... 47
Louisiana..... 47
Waxwing, Cedar..... 41
Whip-poor-will..... 30
Woodcock, American..... 21
Woodpecker, Downy..... 29
Hairy..... 29
Red-bellied..... 30
Red-headed 30
Wren, Carolina..... 49
House..... 49
Long-billed Marsh..... 50
Winter..... 49
Yellow-legs..... 23
Greater..... 22
Yellow-throat, Maryland..... 47

Digitized by Google

PRESS OF THE
ST. JOHNSBURY REPUBLICAN CO.,
St. Johnsbury, Vt.

Digitized by Google

Digitized by Google

Digitized by Google

Digitized by Google

Digitized by Google

Digitized by Google

3 2044 107 163 032

Digitized by Google

Digitized by Google

3 20



Digitized by Google

Digitized by Google

Digitized by Google